


1  
2  
3  
4

**An-Najah N. University**

**Faculty of Graduate Studies**



**THE S-PROPERTY AND BEST APPROXIMATION**

**By : Sawsan Azmi Sabri AL-Dwaik**

**Supervisors : Dr. Abdallah A. Hakawati**

**Dr. Waleed Deeb**

**Submitted In Partial Fulfillment of the Requirements  
For The Degree of Master of Mathematics , Faculty of  
Graduate Studies , At An-Najah National University At  
Nablus , Palestine .**

**2000**

# THE S-PROPERTY AND BEST APPROXIMATION


By : Sawsan Azmi Sabri AL-Dwaik

This Thesis was defended successfully on - August - 2000 and approved by


## Committee Members

## Signature


1. Dr. Abdallah A. Hakawati . (An-Najah University)



2- Dr. Waleed Deeb . (The Arab American University)




3. Dr. Mahmoud Masri . (An-Najah University)



4. Dr. Bassam A. Manasrah . (Hebron University)





## الإهداء

إلى الذين فضلهم بلغ السماء  
وسيظل رضاهم موضع الرجاء  
إليكم والدي ووالدتي وأخوتي  
حبا وعهداً مني بالوفاء

## **Acknowledgments**

First of all , I thank my God for all the blessing he bestowed on me and continues to bestow on me .

I would like to express my gratitude and appreciation for my brother Azam AL-Dwaik for his help , patience , encouragement , supporting this project , and for his excellent job in typing the first copy .

My sincere thanks go to my advisor Dr. Abdallah A. Hakawati who offered me advice and assistance of every stage of my thesis including wise supervision and constant encouragement . His fruitful suggestions were very helpful .

I am indebted to Dr. W. Deeb who spared his valuable time to go through the proofs and made many worthwhile suggestion . I thank Dr. Deeb for his valuable and constructive comments which substantially improved the final product and for the time and effort he put .

I thank my kind , sympathetic mother who prays and asks my God to help me . I ask my God to accept her prayer which will benefit me in this life and after death .

I would also like to acknowledge to all my teachers in An-Najah university , department of mathematics .

My thanks are extended to Dr. Bassam A. Manasrah and Mahmoud Masri for serving in my committee .

I thank my family members for the support and encouragement and for their effective cooperation and great care in preparing this thesis .

I thank Dr. Mohammed Abu Eideh (Hebron University) who provided some references and Dr. Maher Qarawani .

Finally it is pleasure to record my thanks to all my friends , and to all employees in Al-Shari'ya school for their spiritual support .

# Contents

	<b>Preface</b>	1
<b>1</b>	<b>Preliminaries</b>	4
	1. Normed Linear Spaces	4
	2. Compactness	8
	3. Banach Space and Hilbert Space	9
	4. Projection	11
	5. Lebesgue Measure and Integrable Function	12
<b>2</b>	<b>Approximation in Normed Spaces</b>	15
	1. Introduction	15
	2. The Set of Best Approximants	15
	3. Proximinal Set	19
	4. Approximation in Hilbert Space	25
<b>3</b>	<b>Best Approximation in Orlicz Space</b>	28
	1. Orlicz space	28
	2. Best Approximation in $L^\phi(\mu, X)$	31
<b>4</b>	<b>S-Property</b>	42
	1. Introduction	42
	2. Further Results	50
	<b>References</b>	60

542787

## Preface

The problem of best approximation is the problem of finding, for a given point  $x$  and a given set  $G$  in a normed space  $(X, \|\cdot\|)$ , a point  $g_0$  in  $G$  which should be nearest to  $x$  among all points of the set  $G$ .

However, in our study, we shall mainly take as  $X$  not an arbitrary normed space but Orlicz space, we shall denote by  $P(x, G)$ , the set of all elements of best approximants of  $x$  in  $G$ .

$$\text{i.e } P(x, G) = \{ g_0 \in G : \|x - g_0\| = \inf \{ \|x - g\| : g \in G \} \}$$

The problem of best approximation began, in 1853, with P. L. Chebyshev who considered the problem in the space of all real valued continuous function defined on  $[a, b]$ , a closed real interval in  $\mathbb{R}$ .

My thesis consist of four chapters. Each chapter is divided into sections. A number like 2.1.3 indicates item (definition, theorem, corollary or lemma) number 3 in section 1 of chapter 2. Each chapter begins with a clear statement of the pertinent definitions and theorems together with illustrative and descriptive material. At the end of this thesis we present a collection of references.

In chapter (1) we introduce the basic results and definitions which shall be needed in the following chapters. The topics include projection, normed space, compactness, Hilbert space and measure theory. This chapter is absolutely fundamental. The results have been stated without proofs, for

theory may be looked up in any standard text book in Functional Analysis .  
 A reader who is familiar with these topics may skip this chapter and refer to it only when necessary .

Chapter (2) will be devoted to give an introduction to fundamental ideas of Best Approximation in Normed Space . We will start by introducing the definition of best approximants of  $x \in X$  in a closed subspace  $G$  of  $X$  . We denote the set of all best approximation of  $x$  in  $G$  by  $P(x,G)$  . In section (2) we study the properties of  $P(x,G)$  . In section (3) we define proximal set and Chebyshev subspace , and we mention some conditions that can assure that  $G$  is proximal in  $X$  . Finally , we define  $L^p$ -summand and give a simpler proof for the fact that “every a closed subspace of a Hilbert space is proximal ” .

Chapter (3) has two purposes . First , we review the properties of Orlicz spaces . Second , we introduce some of the basic theory of proximality in Orlicz space . This material was designed to meet the needs of chapter (4) .

W. Deeb and R. Khalil proved the following results .

- (1) If  $G$  is 1-complemented in  $X$  , then  $G$  is proximal in  $X$  . [1 , p.529] .
- (2) If  $L^\phi(\mu,G)$  is proximal in  $L^\phi(\mu,X)$  , then  $G$  is proximal in  $X$  . [3 , p.8] , [2 , p.297] , [4 , p.37] .
- (3) If  $L^1(\mu,G)$  is proximal in  $L^1(\mu,X)$  , then  $L^\infty(\mu,G)$  is proximal in  $L^\infty(\mu,X)$  . [1 , p.528] .



Some questions about proximality in  $L^\phi(\mu, X)$  now suggest themselves .

(1) Let  $X$  be a Banach space and let  $G$  be proximal in  $X$  . Under what conditions can it be asserted that  $G$  is 1-complemented in  $X$  ?

(2) If  $G$  is proximal in  $X$  , Under what conditions can it be concluded that  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  ? In particular , is the proximality of  $G$  in  $X$  a sufficient condition ?

(3) If  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$  . Under what condition can be asserted that  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  .

These questions are addressed in the section (1) of chapter (4) .

The answer depends on the S-property .

Some interesting results have been achieved . Among of which it is shown that if  $G$  has the S-property then  $L^\phi(\mu, G)$  has the S-property . It is also proved that if  $G$  has the S-property then

$$L^\phi(\mu, P_G^{-1}(0)) = P_{L^\phi(\mu, G)}^{-1}(0) .$$

I ask our God to be our assistant to continue our efforts so as to achieve the hopes and desires of all scholars in mathematics .

## Chapter 1

### Preliminaries

This chapter contains some definitions and basic result about normed space , Hilbert space , compactness , Banach space , projection and measure theory which will be used in the subsequent chapters .

#### 1. Normed Linear Spaces :

**Definition 1.1.1 :** [9 , p.35] . Let  $X$  be a linear space over  $K$  . A norm on  $X$  is a function  $\|\cdot\| : X \rightarrow \mathbf{R}$  such that for  $x, y \in X$  and  $k \in K$  , we have :

(i)  $\|x\| \geq 0$  and  $\|x\| = 0$  if and only if  $x = 0$  .

(ii)  $\|x + y\| \leq \|x\| + \|y\|$  .

(iii)  $\|kx\| = |k| \|x\|$  .

A normed linear space  $X$  is a linear space  $X$  with a norm  $\|\cdot\|$  on it .

**Theorem 1.1.2 :** [9 , p.35] .

(1) Every normed space is a metric space with respect to the metric

$$d(x, y) = \|x - y\| .$$

(2) For any two elements  $x$  and  $y$  of a normed space we have ,

$$| \|x\| - \|y\| | \leq \|x - y\| .$$

(3) A norm is a real-valued continuous function .

**Definition 1.1.3 :** [9 , p.18] . Let  $(X, \|\cdot\|)$  be a normed linear space and  $G \subset X$  . For  $x \in X$  , we define the distance of a point  $x$  from the subset  $G$

as :

$$d(x,G) = \inf \{ \|x - y\| : y \in G \}$$

**Theorem 1.1.4** : [ 12 , p.147] . Let  $X$  be a normed linear space and  $G$  be a subspace of  $X$ , then

- (1)  $d(x + g, G) = d(x, G)$  ( $x \in X, g \in G$ )  
 (2)  $d(x + y, G) \leq d(x, G) + d(y, G)$  ( $x, y \in X$ )  
 (3)  $d(\alpha x, G) = |\alpha| d(x, G)$  ( $x \in X, \alpha = \text{scalar}$ )  
 (4)  $|d(x, G) - d(y, G)| \leq \|x - y\|$  ( $x, y \in X$ )

**Proof** : For (1) . Let  $x \in X, g \in G$  and  $\epsilon > 0$  be arbitrary . By the definition of  $d(x, G) = \inf \{ \|x - g\| : g \in G \}$  there exist  $g_0 \in G$  s.t

$$\|x - g_0\| \leq d(x, G) + \epsilon$$

consequently , we have

$$d(x + g, G) \leq \|x + g - (g_0 + g)\| = \|x - g_0\| \leq d(x, G) + \epsilon$$

But  $x \in X, g \in G$  and  $\epsilon > 0$  were arbitrary , hence

$$d(x + g, G) \leq d(x, G) \quad (x \in X, g \in G) \dots\dots(1)$$

Applying these relation for  $x + g \in X$  instead of  $x$  and  $-g \in G$  instead of  $g \in G$ , we obtain

$$d(x, G) \leq d(x + g, G) \quad (x \in X, g \in G) \dots\dots(2)$$

From (1), (2) we get  $d(x + g, G) = d(x, G)$  ( $x \in X, g \in G$ ) .

For (2) of the theorem ; Let  $x, y \in X$  and  $\epsilon > 0$  be arbitrary . By the definition of  $d(x, G)$  and  $d(y, G)$  there exist  $g_1, g_2 \in G$  s.t

$$\|x - g_1\| \leq d(x, G) + \epsilon / 2, \quad \|y - g_2\| \leq d(y, G) + \epsilon / 2.$$

Consequently, we have

$$d(x+y, G) \leq \|x + y - (g_1 + g_2)\| \leq \|x - g_1\| + \|y - g_2\| \leq d(x, G) + d(y, G) + \epsilon$$

But  $x, y \in X$ , and  $\epsilon > 0$  were arbitrary, hence

$$d(x + y, G) \leq d(x, G) + d(y, G) \quad (x, y \in X).$$

For (3) : Let  $x \in X$ ,  $\alpha \neq 0$  be a scalar and  $\epsilon > 0$  be arbitrary and take  $g_0 \in G$  satisfying

$$\|x - g_0\| \leq d(x, G) + \epsilon / |\alpha|$$

We have

$$d(\alpha x, G) \leq \|\alpha x - \alpha g_0\| = |\alpha| \|x - g_0\| \leq |\alpha| d(x, G) + \epsilon$$

But  $x, \alpha \neq 0$  and  $\epsilon$  were arbitrary, it follows that

$$d(\alpha x, G) \leq |\alpha| d(x, G) \dots\dots\dots(1)$$

Applying this relation for  $\alpha x$  instead of  $x$  and  $1/\alpha$  instead of  $\alpha$  we obtain

$$d(x, G) = d(1/\alpha \cdot \alpha x, G) \leq 1/|\alpha| d(\alpha x, G)$$

Hence  $|\alpha| d(x, G) \leq d(\alpha x, G) \dots\dots\dots(2)$

From (1) and (2) and since  $d(0, G) = 0$  we get

$$d(\alpha x, G) = |\alpha| d(x, G)$$

For (4) : Let  $x, y \in X$  and  $\epsilon > 0$  be arbitrary and take  $g_0 \in G$  satisfying

$$\|y - g_0\| \leq d(y, G) + \epsilon$$

We have

$$d(x,G) \leq \|x - g_0\| \leq \|x - y\| + \|y - g_0\| \leq \|x - y\| + d(y,G) + \epsilon$$

But  $x, y$  and  $\epsilon$  were arbitrary, there follows

$$d(x,G) - d(y,G) \leq \|x - y\| \quad (x, y \in X)$$

In these relations, interchange  $x$  and  $y$  yields ;

$$d(y,G) - d(x,G) \leq \|x - y\|$$

Hence  $|d(x,G) - d(y,G)| \leq \|x - y\|$ . ■

**Definition 1.1.5 :** [11 , p.153] .

1) A subset  $H$  of a vector space  $X$  is called a hyperplane if there exists a linear functional  $f \neq 0$  defined on  $X$  such that

$$H = \{ x \in X : f(x) = 0 \}$$

2) A subset  $H$  of a vector space  $X$  is called an affine hyperplane if there exists a linear functional  $f \neq 0$  defined on  $X$  and a real number  $\alpha$  such that

$$H = \{ x \in X : f(x) = \alpha \} .$$

**Theorem 1.1.6 :** [12 , p.24] . Let  $X$  be a normed linear space, and  $H = \{ y \in X, f(y) = \alpha \}$  be a hyperplane of  $X$ ,  $f$  being a continuous linear functional on  $X$ ,  $\alpha$  a scalar and let  $x \in X$ . Then the distance of the point  $x$  to the hyperplane  $H$  is

$$d(x,H) = |f(x) - \alpha| / \|f\|$$

**Theorem 1.1.7 :** [7 , p.74] . Every finite dimensional subspace  $G$  of a normed space  $X$  is closed in  $X$ .

Now for linear maps , we have the following theorem .

**Theorem 1.1.8 :** [11 , p.26] . Let  $X$  and  $Y$  be normed spaces .

(a) A linear map  $T : X \rightarrow Y$  is continuous if and only if  $T$  is bounded .

(b) The null space  $N(T)$  of a non zero continuous linear map is a closed subspace of  $X$  .

## 2. Compactness :-

**Definition 1.2.1 :** [7 , p.77] . A metric space  $X$  is said to be compact if every sequence in  $X$  has a convergent subsequence . A subset  $M$  of  $X$  is said to be compact if every sequence in  $M$  has a convergent subsequence whose limit is an element of  $M$  .

A general property of compact sets is expressed in :

**Lemma 1.2.2 :** [7 , p.77] . A compact subset of a metric space is closed and bounded .

However , for a finite dimensional normed space we have :

**Theorem 1.2.3 :** [7 , p.77] . In a finite dimensional normed space  $X$  , a subset of  $X$  is compact if and only if it is closed and bounded .

In connection with continuous mapping a fundamental property is that compact sets have compact images , as follow we have :

**Theorem 1.2.4 :** [7 , p.81] . Let  $X$  and  $Y$  be metric spaces and  $T : X \rightarrow Y$  be a continuous mapping . Then the image of a compact subset  $M$  of  $X$  is compact .

From this theorem we conclude that the following property, well-known from Calculus for continuous functions, carries over to metric space.

**Theorem 1.2.5** : [7, p.81]. A continuous mapping  $T$  of a compact subset  $M$  of a metric space  $X$  into  $\mathbb{R}$  assumes its maximum and minimum on  $M$ .

### 3. Banach Space and Hilbert Space :-

**Definition 1.3.1** : [7, p.58]. A complete normed space is called a Banach space.

**Remark 1.3.2** : [9, p.47].

(1) The set of all bounded linear maps on a normed space  $X$  into a normed space  $Y$  is denoted by  $B[X, Y]$ . If  $X = Y$ ,  $B[X]$  denotes  $B[X, X]$ .

(2) The set of all bounded linear functionals on a normed space  $X$  is denoted by  $X^*$ .

For  $f \in X^*$ , any element  $x \in X - \{0\}$  with the property that

$$f(x) = \|f\| \|x\| \text{ is called a maximal element of } f.$$

**Definition 1.3.3** : [12, p.110]. A normed space  $X$  is strictly convex if and only if every functional  $f \neq 0 \in X^*$  has at most one maximal element of norm 1. An equivalent form : a normed space  $X$  is strictly convex if

$$\|x + y\| = \|x\| + \|y\| \text{ and } \|x\| = \|y\| \text{ implies that } x = y.$$

**Remark 1.3.4** : [11, p.32]. An arbitrary  $f \in c^*$  can be expressed as

$$f(x) = y_0 \lim_{n \rightarrow \infty} x_n + \sum_{n=1}^{\infty} y_n x_n \quad \text{where}$$

$x = (x_1, x_2, x_3, \dots) \in \mathbb{C}$  and  $y = (y_0, y_1, y_2, \dots)$  such that  $\sum_{n=1}^{\infty} |y_n| < \infty$  and

$$\|f\| = |y_0| + \sum_{n=1}^{\infty} |y_n|$$

**Definition 1.3.5 :** [9 , p.176] . Let  $X$  be a vector space over the field  $K$  . An inner product on  $X$  is a function

$\langle \cdot, \cdot \rangle : X \times X \rightarrow K$  such that for all  $x, x', y \in X$  and  $\alpha \in K$  .

$$(1) \langle x + x', y \rangle = \langle x, y \rangle + \langle x', y \rangle$$

$$(2) \langle \alpha x, y \rangle = \alpha \langle x, y \rangle$$

$$(3) \overline{\langle x, y \rangle} = \langle y, x \rangle$$

$$(4) \langle x, x \rangle \geq 0 \quad \forall x \in X \quad \text{and} \quad \langle x, x \rangle = 0 \quad \text{iff} \quad x = 0 .$$

An inner product space is a linear space with an inner product on it .

**Definition 1.3.6 :** [9 , p.182] . A Hilbert space  $X$  is a Banach space in which the norm satisfies the parallelogram law .

$$\|x + y\|^2 + \|x - y\|^2 = 2 \|x\|^2 + 2 \|y\|^2, \quad x, y \in X$$

**Definition 1.3.7 :** [7 , p.131] . Two vectors  $x$  and  $y$  in an inner-product space are called orthogonal , (written  $x \perp y$ ) , if  $\langle x, y \rangle = 0$  .

**Theorem 1.3.8 :** [7 , p.135] . Let  $X$  be an inner-product space and  $x, y \in X$  . Then for  $x \perp y$  we have  $\|x + y\|^2 = \|x\|^2 + \|y\|^2$  .



In the case of a general Hilbert space  $X$ , we obtain an interesting representations of  $X$  as a direct sum of a closed subspace  $M$  and its orthogonal complement

$$M^\perp = \{ x \in X, x \perp M \}$$

which is the set of all vectors orthogonal to each member of  $M$ . The next theorem is sometimes called the projection theorem.

**Theorem 1.3.9** : [11, p.96]. If  $M$  is a closed subspace of a Hilbert space  $X$ , then

$$X = M \oplus M^\perp$$

#### 4. Projection :-

**Definition 1.4.1** : [9, p.85]. If  $X$  is a normed space and  $P \in B[X]$  satisfies  $P^2 = P$ , then  $P$  is called a projection.

**Theorem 1.4.2** : [9, p.85]. If  $P$  is a projection on a Banach space  $X$ , and if  $M$  and  $N$  are its range and null space, respectively, then  $M$  and  $N$  are closed subspaces and

$$X = M \oplus N$$

**Theorem 1.4.3** : [9, p.85]. Let  $X$  be a Banach space, and let  $M$  and  $N$  be closed subspace of  $X$  such that

$$X = M \oplus N$$

The mapping defined on each  $z = x + y, x \in M, y \in N$  by  $P(z) = x$  is a projection on  $X$  whose range is  $M$  and whose null space is  $N$ .

## 5. Lebesgue Measure and Integrable Function :-

In this section we shall review the theory of the Lebesgue measure and the p-integrable function .

**Definition 1.5.1** : [10 , p.9] .

(a) A collection  $\mathcal{M}$  of subsets of a set  $X$  is said to be a  $\sigma$ -algebra of  $X$  if it has the following properties .

(i)  $X \in \mathcal{M}$

(ii) If  $A \in \mathcal{M}$  , then  $A^c \in \mathcal{M}$  where  $A^c$  is the complement of  $A$  relative to  $X$  .

(iii) If  $A = \bigcup_{n=1}^{\infty} A_n$  ,  $A_n \in \mathcal{M}$  for  $n = 1, 2, 3, \dots$  , then  $A \in \mathcal{M}$  .

(b) If  $\mathcal{M}$  is a  $\sigma$ -algebra in  $X$  , then  $X$  is called a measurable space , and the members of  $\mathcal{M}$  are called the measurable sets in  $X$  .

(c) If  $X$  is a measurable space ,  $Y$  is a topological space , and  $f$  is a mapping of  $X$  into  $Y$  , then  $f$  is said to be measurable provided that  $f^{-1}(V)$  is a measurable set in  $X$  for every open set  $V$  in  $Y$  .

For  $E \subset X$  , let  $\chi_E$  denote the characteristic function of  $E$  ; i.e ,

$$\chi_E(t) = \begin{cases} 1 & \text{if } t \in E \\ 0 & \text{if } t \notin E \end{cases}$$

It is measurable iff  $E$  is measurable .

**Definition 1.5.2 :** [8 , p.113] . A function  $f : T \rightarrow X$  is said to be simple if its range contains only finitely many points  $x_1, x_2, \dots, x_n$  and if  $f^{-1}(x_i)$  is measurable for  $i = 1, 2, \dots, n$  . Such a function then can be written as

$$f = \sum_{i=1}^n x_i \chi_{E_i} \text{ where for each } i, E_i = f^{-1}(x_i) . \text{ Define}$$

$$\int_E f d\mu = \sum_{i=1}^n x_i \mu(E_i \cap E)$$

If  $f$  is a non-negative measurable function on  $E$  , define

$$\int_E f d\mu = \sup \left\{ \int_E s d\mu : 0 \leq s \leq f, s \text{ is simple and measurable on } E \right\}$$

**Definition 1.5.3 :** [10 , p.17] .

(a) A measure is a function  $\mu$  , defined on a  $\sigma$ -algebra  $\mathcal{M}$  , whose range is in  $[0, \infty]$  and which is countably additive . This means that if  $\{A_n\}$  is a disjoint countable collection of members of  $\mathcal{M}$  , then

$$\mu \left( \bigcup_{n=1}^{\infty} A_n \right) = \sum_{n=1}^{\infty} \mu (A_n)$$

(b) A measure space is a measurable space which has a measure defined on the  $\sigma$ -algebra of its measurable sets .

A property which is true except for a set of measure zero is said to hold almost everywhere .

**Remark 1.5.4 :** [10 , p.21] . The following propositions are immediate consequences of the definitions . Functions and sets are assumed to be measurable on a measure space  $E$  .

(a) If  $0 \leq f \leq g$  then  $\int_E f \leq \int_E g$ .

(b) If  $A \subset B$  and  $f > 0$ , then  $\int_A f d\mu \leq \int_B f d\mu$ .

(c) If  $c$  is constant, then  $\int_E c d\mu = c\mu(E)$ .

(d) If  $E = E_1 \cup E_2$  where  $E_1$  and  $E_2$  are disjoint, then

$$\int_E f d\mu = \int_{E_1} f d\mu + \int_{E_2} f d\mu$$

(e) If  $f \geq 0$  and  $\int_E f d\mu = 0$ , then  $f = 0$  almost everywhere on  $E$ .

**Definition 1.5.5** : [1, p.527]. Let  $X$  be a real Banach space, and  $(T, \mu)$  be a finite measure space. The space of Bochner  $p$ -integrable functions defined on  $(T, \mu)$  with values in a Banach space  $X$  is denoted  $L^p(\mu, X)$ .

For  $f \in L^p(\mu, X)$ , we write

$$\|f\|_p = \begin{cases} \left( \int_T \|f(t)\|^p d\mu(t) \right)^{1/p} & 1 \leq p < \infty \\ \int_T \|f(t)\|^p d\mu(t) & 0 < p < 1 \\ \text{ess. sup}_{t \in T} \|f(t)\| & p = \infty \end{cases}$$

where  $\text{ess. sup}_{t \in T} \|f(t)\| = \inf \{ M : \mu \{ t : \|f(t)\| > M \} = 0 \}$ .

## Chapter 2

### Approximation in Normed Spaces

**1. Introduction :** Let  $X = (X, \|\cdot\|)$  be a normed space,  $G$  be a subspace of  $X$  and  $x \in X$ , an element  $g_0 \in G$  is called a **best approximant** of  $x$  in  $G$  if

$$\|x - g_0\| = d(x, G) = \inf \{\|x - g\| : g \in G\}.$$

We see that for  $x \in X$  a best approximant  $g_0 \in G$  is an element of minimal distance from the given  $x$ . Such a  $g_0 \in G$  may or may not exist.

We shall denote the set of all elements of best approximants of  $x$  in  $G$  by  $P(x, G)$  i.e  $P(x, G) = \{g \in G : \|x - g\| = d(x, G)\}$ .

#### **2. The Set of Best Approximants :-**

In this section, we introduce some basic properties of  $P(x, G)$ , we use the book of Erwin Kreyszig for this purpose, so we start with the following.

**Example 2.2.1 :** Let  $X = (X, \|\cdot\|_1)$  be the normed space of ordered pairs  $x = (x_1, x_2)$  of real numbers with norm defined by  $\|x\|_1 = |x_1| + |x_2|$ .

Let us take  $x = (1, -1)$  and the subspace  $G = \{g = (g_1, g_1) : g_1 \in \mathbf{R}\}$  then for all  $g \in G$ , we clearly have

$$\|x - g\|_1 = |1 - g_1| + |-1 - g_1| \geq |1 - g_1 + 1 + g_1| = 2.$$

Hence  $d(x, G) \geq 2$  .....(1)

Also  $(1, 1) \in G$ , and  $d(x, G) \leq \|x - (1, 1)\| = 2$  .....(2)

From (1) and (2) we get  $d(x, G) = 2$  and

$$P(x,G) = \{ g \in G : g = (g_1, g_2) \text{ and } |g_1| \leq 1 \} .$$

From this example we conclude that  $P(x,G)$  need not be a subspace .

**Theorem 2.2.2 :** Let  $G$  be a subspace of a normed space  $X$  .

(i) if  $x \in G$  , then  $P(x,G) = \{x\}$  .

(ii) if  $G$  is not closed and  $x \in \overline{G} \setminus G$  , then  $P(x,G) = \phi$  .

**Proof :-** For (i) , let  $x \in G$  , then  $\inf \{ \|x - g\| : g \in G \} = 0$  . Thus , if  $y \in P(x,G)$  then  $\|x - y\| = 0$  hence  $x = y$  since  $X$  is a normed space .

For (ii) , let  $x \in \overline{G} \setminus G$  . This means that  $\forall n \in \mathbf{N} , \exists x_n \in G$

s.t  $\|x_n - x\| \leq 1/n$  . i.e  $d(x,G) = 0$  . Hence  $P(x,G) = \phi$  . ▣

Now provided  $P(x,G) \neq \phi$  , we shall prove that either  $P(x,G)$  contains exactly one element or else an infinite number of elements .

**Theorem 2.2.3 :** Let  $X$  be a normed space ,  $x \in X$  , and let  $G$  be a subspace of  $X$  , then  $P(x,G)$  is a convex set .

**Proof :** Let  $\delta$  be the distance from  $x$  to  $G$  . The statement holds if  $P(x,G)$  is empty or has just one point . Now suppose that  $y , z \in P(x,G)$  such that  $y \neq z$  . So

$$\|x - y\| = \|x - z\| = \delta .$$

We will show that if  $0 \leq \alpha \leq 1$  , and if

$$w = \alpha y + (1 - \alpha)z , \text{ then } w \in P(x,G)$$

to show this :

$$\begin{aligned}
\|x - w\| &= \|x - (\alpha y + (1 - \alpha)z)\| \\
&= \|x - \alpha y - (1 - \alpha)z + \alpha x - \alpha x\| \\
&= \|\alpha(x - y) + (1 - \alpha)(x - z)\| \\
&\leq \alpha \|x - y\| + (1 - \alpha) \|x - z\| \\
&= \alpha \delta + (1 - \alpha)\delta \\
&= \delta
\end{aligned}$$

Therefore  $\|x - w\| \leq \delta \dots\dots\dots(1)$

Also  $w \in G$ , since  $G$  is a subspace, so ;  $\delta \leq \|x - w\| \dots\dots\dots(2)$

From (1) and (2) we get that  $\|x - w\| = \delta$ , so  $w \in P(x, G)$ . Since

$y, z \in P(x, G)$  were arbitrary ;  $P(x, G)$  is convex. ■

**Theorem 2.2.4** : Let  $G$  be a subspace of a normed space  $X$ , then for  $x \in X$ .

(i)  $P(x, G)$  is a bounded set .

(ii) If  $G$  is a closed subspace, then  $P(x, G)$  is a closed set .

**Proof** : For (i), let  $g_0 \in P(x, G)$ , we have by definition  $\|x - g_0\| = \delta$  where

$$\delta = \inf \{ \|x - g\| : g \in G \} .$$

Now  $\|g_0\| = \|g_0 - x + x\|$

$$\leq \|g_0 - x\| + \|x\|$$

$$\leq \|x\| + \|x\| , \quad \text{since } 0 \in G$$

$$= 2 \|x\|$$

Thus  $P(x,G)$  is bounded .

For (ii) , we show that if a sequence  $(g_n) \in P(x,G)$  such that  $g_n \rightarrow g$  then  $g \in P(x,G)$  . Now  $g_n \in P(x,G) \forall n \in \mathbb{N}$  , so  $\|x - g_n\| = d(x,G) = \delta$  ,  $\forall n \in \mathbb{N}$ .

Also  $g_n \in G$  . Since  $G$  is a closed subspace , then  $g \in G$  .

But the function  $F_x : G \rightarrow \mathbb{R}$  defined by  $F_x(g) = \|x - g\| \forall g \in G$  is continuous by part (3) of Theorem (1.1.2) . So  $F_x(g_n) \rightarrow F_x(g)$  implies that

$$\|x - g_n\| \rightarrow \|x - g\|$$

But  $\|x - g_n\| = \delta \forall n \in \mathbb{N}$  , so  $\|x - g\| = \delta$  .

Therefore  $g \in P(x,G)$  . ■

**Theorem 2.2.5 :** Let  $G$  be a subspace of a normed space  $X$  . For  $x \in X$  :

(i) if  $z \in P(x,G)$  then  $\alpha z \in P(\alpha x, G)$  for all scalars  $\alpha$  .

(ii) if  $z \in P(x,G)$  then  $z + g \in P(x + g, G)$  for all  $g \in G$  .

**Proof :** For (i) ; if  $g \in G$  and  $\alpha$  a scalar  $\neq 0$  we have

$$\|\alpha x - g\| = |\alpha| \|x - (1/\alpha).g\| \geq |\alpha| \|x - z\| = \|\alpha x - \alpha z\|$$

Thus  $\alpha z \in P(\alpha x, G)$  .

For (ii) ; if  $g' \in G$  we have

$$\|x + g - g'\| \geq \|x - z\| = \|x + g - (z + g)\|$$

whence  $z + g \in P(x + g, G)$  . ■

An element  $x$  of a normed linear space  $X$  is said to be **orthogonal** to an



element  $y \in X$ , and we write  $x \perp y$ , if we have  $\|x + \alpha y\| \geq \|x\|$  for every scalar  $\alpha$ . In a Hilbert space  $X$  we have  $x \perp y$  if and only if  $\langle x, y \rangle = 0$ .

An element  $x$  of a normed linear space  $X$  is said to be orthogonal to a set  $G \subset X$  and we write  $x \perp G$ , if we have  $x \perp g$  ( $g \in G$ ).

The relationship between orthogonality and best approximation is given by the following theorem.

**Theorem 2.2.6** : [12, p.92]. Let  $X$  be a normed space,  $G$  a subspace of  $X$ ,  $x \in X \setminus \overline{G}$  and  $g_0 \in G$ . We have  $g_0 \in P(x, G)$  if and only if  $x - g_0 \perp G$ .

**Proof** : By the definition, orthogonality means that

$$\|x - g_0 + \alpha g\| \geq \|x - g_0\| \quad (g \in G, \alpha = \text{scalar})$$

and this is obviously equivalent to saying that  $g_0 \in P(x, G)$ . ■

### 3. Proximinal Set :-

Theorem 2.2.2 shows that if  $X$  is a normed space and  $G$  a subspace of  $X$ , then for every  $x \in G$  the set  $P(x, G)$  is non empty, and if the subspace  $G$  is not closed, then for every  $x \in \overline{G} \setminus G$  the set  $P(x, G)$  is empty.

Furthermore, for the elements  $x$  of  $X \setminus \overline{G}$  the set  $P(x, G)$  may or may not be empty. The subspaces  $G \subset X$  which have the property that  $P(x, G) \neq \emptyset$  for every  $x \in X$  are called **proximinal sets in  $X$** . Some authors use the term distance set, or existence set for proximinal sets.

**Remark 2.3.1** : In a normed spaces  $X$ , the condition that  $G$  is a closed

subspace of  $X$  is not sufficient for  $G$  to be proximal in  $X$  as shown by the following example .

**Example :** Let  $X = c_0$  = the space of all sequences of complex numbers that converge to zero , with norm  $\|x\| = \sup_n |x_n|$  , and let

$$G = \{ x = (x_n) \in c_0 : \sum_{n=1}^{\infty} 2^{-n} x_n = 0 \} .$$

$G$  is closed but not proximal . To show that define a linear functional

$$f : c_0 \rightarrow \mathbb{C} \text{ by } f(x) = \sum_{n=1}^{\infty} 2^{-n} x_n \text{ for all } x = (x_n) \in c_0 .$$

Then  $f \in c_0^*$  and  $\|f\| = 1$  by Remark (1.3.4) and hence  $G$  is a closed subspace of  $c_0$  by part (2) of Theorem (1.1.8) .

Now let  $x = e^{(1)} = (1, 0, 0, \dots) \in c_0$  .

Then  $d(e^{(1)}, G) = 1/2$  by Theorem (1.1.6) .

We claim that there does not exist any  $g \in G$  s.t  $\|e^{(1)} - g\| = 1/2$  .

Assume on the contrary that  $\exists g = (g_i) \in G$  s.t  $\|e^{(1)} - g\| = 1/2$  , then

$$|1 - g_1| \leq 1/2 \text{ and}$$

$$|g_k| \leq 1/2 \text{ for all } k \geq 2 . \text{ Since } \sum_{n=1}^{\infty} 2^{-n} g_n = 0 , \text{ we get that}$$

$$1/4 \leq 1/2 |g_1| = \left| \sum_{n=2}^{\infty} 2^{-n} g_n \right| \leq \sum_{n=2}^{\infty} 2^{-n} |g_n| \leq 1/2 \sum_{n=2}^{\infty} 2^{-n} = 1/4 \dots\dots\dots *$$

So we must have equalities in (\*), and that can happen only if  $|g_n| = 1/2$  for all  $n$  . But this contradicts our assumption that  $g \in c_0$  . Thus  $G$  is not proximal in  $c_0$  . ■

We call a linear subspace  $G$  of a normed space  $X$  a **semi-Chebyshev** subspace if for every  $x \in X$  the set  $P(x,G)$  contains at most one element .

An example of such subspaces is that of the subspaces  $G$  with the property that the set  $P(x,G)$  is empty for all  $x \in X \setminus G$  . We will see such a space in Remark (4.1.2) .  $G$  is called a **Chebyshev subspace** if it is simultaneously proximal and semi-Chebyshev . i.e if for every  $x \in X$  the set  $P(x,G)$  contains exactly one element .

**Theorem 2.3.2** : For a subspace  $G$  of a normed space  $X$  , the following are equivalent :

(i)  $G$  is proximal in  $X$  .

(ii)  $X = G + P_G^{-1}(0)$  where  $P_G^{-1}(0) = \{ x \in X : 0 \in P(x,G) \}$  .

**Proof** : If  $G$  is proximal and  $x \in X$  ,  $g_0 \in P(x,G)$  , then

$$x = g_0 + (x - g_0) \in G + P_G^{-1}(0) .$$

Conversely if we have (ii) and  $x \in X$  ,  $x = g_0 + y$  where  $g_0 \in G$  ,  $y \in P_G^{-1}(0)$  then  $0 \in P(y,G) = P(x - g_0,G)$  , implies  $d(x - g_0,G) = \|x - g_0\|$  .

$$\Rightarrow d(x,G) = \|x - g_0\| . \text{ Hence } g_0 \in P(x,G) . \quad \blacksquare$$

**Theorem 2.3.3** : Let  $G$  be a compact subspace of a normed space  $X$  , then  $G$  is proximal in  $X$  .

542787

**Proof** : Fix  $x \in X$  , since the mapping  $T_x : G \rightarrow \mathbb{R}$  defined by  $T_x(g) = \|x - g\|$  is continuous by part (3) of Theorem (1.1.2) , then  $T(G) \subset \mathbb{R}$  is compact by

Theorem (1.2.4). So  $T(G)$  is closed and bounded in  $\mathbf{R}$  by Theorem (1.2.3) .

Theorem (1.2.5) implies  $\inf T(G) \in T(G) = \{ \|x - g\| : g \in G \}$  . Hence there exists  $g_0 \in G$  such that  $\|x - g_0\| = \inf \{ \|x - g\| : g \in G \}$  . This proves that  $g_0 \in P(x,G)$  . Since  $x$  is an arbitrary ,  $P(x,G) \neq \phi$  for all  $x \in X$  . ■

While compactness of  $G$  is a sufficient condition for a best approximant to exist , it is clearly not necessary . For , consider the noncompact subspace  $G = \{ (x,0) : x \in \mathbf{R} \}$  of  $\mathbf{R}^2$  . Clearly  $G$  is proximal in  $\mathbf{R}^2$  .

**Corollary 2.3.4** : Every closed and bounded subspace of a finite dimensional normed space  $X$  is proximal in  $X$  .

**Proof** : Every closed and bounded subspace of a finite dimensional space is compact by Theorem (1.2.3) . ■

**Theorem 2.3.5** : Let  $G$  be a finite dimensional subspace of a normed space  $X$  , then  $G$  is proximal in  $X$  .

**Proof** : Let  $x \in X$  be given , consider the closed Ball .

$$\bar{B} = \{ g \in G : \|g\| \leq 2 \|x\| \}$$

Then  $0 \in \bar{B}$  , so that for the distance from  $x$  to  $\bar{B}$  we obtain

$$d(x, \bar{B}) = \inf \{ \|x - g'\| : g' \in \bar{B} \} \leq \|x - 0\| = \|x\|$$

Now if  $g \in G$  and  $g \notin \bar{B}$  , then  $\|g\| > 2 \|x\|$  and

$$\|x - g\| \geq \|g\| - \|x\| > \|x\| \geq d(x, \bar{B}) \dots\dots\dots(1)$$

If  $g \in G$  and  $g \in \overline{B}$ , then  $d(x, \overline{B}) \leq \|x - g\| \dots\dots\dots(2)$

(1) and (2) imply that  $d(x, \overline{B})$  is a lower bound of the set

$$\{ \|x - g\| : g \in G \}.$$

Hence  $d(x, \overline{B}) \leq d(x, G)$ .

Also  $\overline{B} \subset G \Rightarrow d(x, G) \leq d(x, \overline{B})$ .

This shows that  $d(x, \overline{B}) = d(x, G)$ . Since  $\overline{B}$  is closed and bounded and since  $G$  is finite dimensional; Theorem (1.2.3) implies  $\overline{B}$  is compact.

Theorem (2.3.3) implies that  $\overline{B}$  is proximal in  $X$ . Hence, if a best approximant to  $x$  exists, it must lie in  $\overline{B}$ . Thus  $G$  is proximal in  $X$ . ■

It is not possible to drop the finite dimensional requirement of this theorem. For, let  $X$  be the space of continuous functions defined on  $[0, 1/2]$  with  $L^\infty$  norm. i.e.  $\|f\|_\infty = \max_{0 \leq x \leq 1/2} |f(x)|$ .

Let  $G$  be the subspace of polynomials, and let  $g(x) = 1/(1-x)$ . For any  $\epsilon > 0$ , there exists a polynomial  $p_\epsilon$  such that  $|g(x) - p_\epsilon(x)| < \epsilon$  for all  $x \in [0, 1/2]$  [ Weierstrass Approximation Theorem ].

Hence  $d(g, G) = 0$ . However, since  $g$  is not a polynomial, we see that there is no  $p \in G$  satisfying  $d(g, G) = \|g - p\| = 0$ .

We present the following example to show that proximal subspaces need

not be finite dimensional ; thus proving that the converse of the foregoing theorem is not true .

**Example 2.3.6 :** The infinite dimensional subspace  $c_0$  of  $c$  is proximal in  $c$  .

**Proof :** On  $c$  , define the linear functional  $f$  by  $f(x) = \lim x_n = \mathbf{x}$  .

Then  $c_0 = \{ x = (x_n) \in c : f(x) = 0 \}$  is the hyperplane of  $c$  and  $d(x,c) = |\mathbf{x}|$  by Theorem (1.1.6) .

Let  $g = (g_n)$  be defined as ,  $g_n = x_n - \mathbf{x}$  .

$$\begin{aligned} \text{Now } g \in c_0 \text{ and } \|x - g\| &= \sup \{ |x_n - g_n| : n \in \mathbf{N} \} \\ &= \sup \{ |x_n - (x_n - \mathbf{x})| : n \in \mathbf{N} \} \\ &= |\mathbf{x}| \end{aligned}$$

Hence  $d(x,c) = \|x - g\|$  and so ;  $g \in P(x,c)$  .

Since  $x$  was arbitrary ;  $c_0$  is proximal in  $c$  . ■

**Theorem 2.3.7:** [12 , p.93] . Let  $X$  be a normed space and  $G$  a hyperplane in  $X$  , passing through  $0$  .  $G$  is proximal in  $X$  if and only if there exists an element  $z \in X - \{0\}$  such that  $0 \in P(z,G)$  . (i.e  $z \perp G$ ) .

**Proof :** Assume that  $G$  is proximal and take arbitrary  $x \in X - G$  ,  $y_0 \in P(x,G)$  . Then for  $z = x - y_0 \neq 0$  we have  $0 \in P(z,G)$  .

Conversely , assume that there exists  $z \in X - \{0\}$  such that  $0 \in P(z,G)$  and let  $x \in X - G$  be arbitrary .

Take  $f \in X^*$  such that

$$G = \{ y \in X : f(y) = 0 \},$$

and put

$$y_0 = x - (f(x) / f(z)) z$$

(we have  $f(z) \neq 0$  since, otherwise  $z \in G$ , so  $0 \in P(z, G) = \{z\}$ , and hence  $z = 0$ , a contradiction to the hypothesis).

We have then

$$f(y_0) = 0$$

Whence  $y_0 \in G$ . Also, since  $(f(z) / f(x)) (y - y_0) \in G$  for every  $y \in G$ , we have

$$\|x - y_0\| = |f(x) / f(z)| \|z\| \leq |f(x) / f(z)| \|z - (f(z) / f(x)) (y - y_0)\| = \|x - y\|$$

Whence  $y_0 \in P(x, G)$ . Since  $x \in X - G$  has been arbitrary, it follows that  $G$  is proximal in  $X$ . ■

#### 4. Approximation in Hilbert space

Before we prove that all closed subspaces of a Hilbert space are proximal we need the following definition.

**Definition 2.4.1 :** [6, p.279]. A closed subspace  $G$  of a Banach space  $X$  is called an  $L^p$ -summand,  $1 \leq p < \infty$ , if there is a bounded projection  $P : X \rightarrow G$  which is onto, and  $\|x\|^p = \|P(x)\|^p + \|x - P(x)\|^p$  for all  $x \in X$ .

**Theorem 2.4.2 :** If  $G$  is an  $L^p$ -summand of a Banach space  $X$ , then  $G$  is proximal in  $X$ .  $1 \leq p < \infty$ .

**Proof :** Let  $x \in X$ , for every  $g \in G$  we have

$$\begin{aligned} \|x - g\|^p &= \|P(x - g)\|^p + \|x - g - P(x - g)\|^p \\ &= \|P(x) - g\|^p + \|x - P(x)\|^p \\ &\geq \|x - P(x)\|^p \end{aligned}$$

Hence  $\|x - g\| \geq \|x - P(x)\|$

i.e  $P(x) \in P(x, G)$ . Thus  $G$  is proximal in  $X$ . ▮

Erwin kreyszig [7] proved that if  $G$  is a closed subspace of a Hilbert space  $X$ , then  $G$  is proximal in  $X$ . He used Cauchy sequence and parallelogram law to prove this theorem. Here we give a simpler proof.

**Theorem 2.4.3 :** Let  $G$  be any closed subspace of a Hilbert space  $X$ , then  $G$  is a Chebyshev subspace.

**Proof :** Since  $G$  is a closed subspace of a Hilbert space  $X$ . Theorem (1.3.9) implies  $X = G \oplus G^\perp$  where  $G^\perp = \{ g \in X : g \perp G \}$ . Hence every element  $x \in X$  has a unique representation i.e  $x = g + z$  where  $g \in G$ ,  $z \in G^\perp$ . Now we define the projection  $P : X \rightarrow G$  by  $P(x) = g$ . Clearly  $P$  is onto and bounded.

Also if  $x = g + z$  and  $z \perp g$  then  $\|z + g\|^2 = \|z\|^2 + \|g\|^2$  by Theorem (1.3.8), so  $\|x\|^2 = \|x - P(x)\|^2 + \|P(x)\|^2$  i.e  $G$  is an  $L^2$ -summand of  $X$ , hence Theorem (2.4.2) implies that  $G$  is proximal in  $X$ .

Now we show that  $P(x, G)$  contains exactly one element.



Let  $x \in X$ , assume  $g_1, g_2 \in P(x, G)$  such that  $g_1 \neq g_2$ . Since  $g_1, g_2 \in G$  and  $G$  is subspace, then  $(g_1 + g_2)/2 \in G$ . By the parallelogram law for :

$(x - g_1)/2$  and  $(x - g_2)/2$  we have

$$\|(x - g_1)/2 + (x - g_2)/2\|^2 + \|(x - g_1)/2 - (x - g_2)/2\|^2 = 2 \|(x - g_1)/2\|^2 + 2 \|(x - g_2)/2\|^2$$

or  $\|x - (g_1 + g_2)/2\|^2 < 1/2 \|x - g_1\|^2 + 1/2 \|x - g_2\|^2 = [d(x, G)]^2$ . This implies that  $\|x - (g_1 + g_2)/2\| < d(x, G)$ .

Which contradicts the definition of  $d(x, G)$ . Hence  $G$  is a Chebyshev subspace. ■

**Corollary 2.4.4 :** Any closed subspace of  $\mathbb{R}^n$ ,  $\mathbb{C}^n$  is a Chebyshev subspace.

## Chapter 3

### Best Approximation in Orlicz Space

#### 1. Orlicz Spaces :

In order to study Orlicz spaces , it is necessary to introduce the definition of modulus functions .

**Definition 3.1.1 :** [5 , p.159] . A function  $\phi : [0, \infty) \rightarrow [0, \infty)$  is called a modulus function if the following are satisfied :

- (i)  $\phi$  is continuous at zero from the right and strictly increasing .
- (ii)  $\phi(0) = 0$  .
- (iii)  $\phi$  is subadditive i.e  $\phi(x + y) \leq \phi(x) + \phi(y)$  .  $\forall x, y \in [0, \infty)$

Examples of such functions are  $\phi(x) = x^p$  ,  $0 < p \leq 1$  , and  $\phi(x) = \ln(1+x)$  .

In fact , if  $\phi$  is a modulus function and  $a \geq 0$  then  $\Psi_1(x) = \phi(x) / (1+\phi(x))$  and  $\Psi_2(x) = a\phi(x)$  are modulus functions . Further , the composition of two modulus functions is a modulus function .

**Theorem 3.1.2 :** Every modulus function is continuous on  $[0, \infty)$  .

**Proof :** Let  $x_0 \in [0, \infty)$  . We show that  $\phi$  is continuous at  $x_0$  , i.e

$\lim_{x \rightarrow x_0} \phi(x) = \phi(x_0)$  . At first we show that

$$|\phi(x) - \phi(y)| \leq \phi(|x - y|) \text{ for all } x, y \in [0, \infty) .$$

Now  $|x| = |x - y + y| \leq |x - y| + |y|$  , since  $\phi$  is strictly increasing and subadditive we get  $\phi(|x|) \leq \phi(|x - y|) + \phi(|y|)$  .

So  $\phi(|x|) - \phi(|y|) \leq \phi(|x - y|)$  .....(1)

Also

$|y| = |y - x + x| \leq |x - y| + |x|$ , implies  $\phi(|y|) - \phi(|x|) \leq \phi(|x - y|)$  .....(2)

From (1) and (2) we get :

$$|\phi(|x|) - \phi(|y|)| \leq \phi(|x - y|) \quad \text{for all } x, y \in [0, \infty).$$

Now given  $\epsilon > 0$ , there exist  $\delta(\epsilon) > 0$  such that if  $0 < x < \delta(\epsilon)$  then  $|\phi(x)| < \epsilon$ , because  $\phi$  is continuous at 0. But  $|\phi(x) - \phi(x_0)| \leq \phi(|x - x_0|) < \epsilon$  if  $|x - x_0| < \delta(\epsilon)$ . Hence  $\phi$  is continuous at  $x_0$ . Since  $x_0$  is arbitrary,  $\phi$  is continuous on  $[0, \infty)$ . ■

**Definition 3.1.3 :** [5, p.159]. Let  $X$  be a real Banach space, and  $(T, \mu)$  be finite measure space. For a given modulus function  $\phi$ , we define the Orlicz space as :

$$L^\phi(\mu, X) = \left\{ f : T \rightarrow X : \int_T \phi(\|f(t)\|) d\mu(t) < \infty \right\}.$$

The function  $d : L^\phi(\mu, X) \times L^\phi(\mu, X) \rightarrow [0, \infty)$ , given by

$$d(f, g) = \int_T \phi(\|f(t) - g(t)\|) d\mu(t)$$

defines a metric on  $L^\phi(\mu, X)$ , under which it becomes a complete metric linear space [4, p.70].

For  $f \in L^\phi(\mu, X)$ , we write

$$\|f\|_\phi = \int_T \phi(\|f(t)\|) d\mu(t).$$

The spaces of  $p$ -Bochner integrable functions  $L^p(\mu, X)$ ,  $0 < p \leq 1$ , by Definition (1.5.5) are reduced to be a special case of the Orlicz space under the modulus function  $\phi(x) = x^p$   $0 < p \leq 1$ .

The following theorem relates  $L^1(\mu, X)$  and  $L^\phi(\mu, X)$  for all modulus functions  $\phi$ .

**Theorem 3.1.4 :** [5, p.159]. If  $\phi$  is a modulus function, then

$$L^1(\mu, X) \subset L^\phi(\mu, X).$$

**Proof :** For each real number  $x$ , we have  $[x] \leq x < [x] + 1$ ; where  $[ ]$  denotes the greatest integer function. But  $\phi$  is strictly increasing and subadditive, then

$$\begin{aligned} \phi(x) &\leq \phi([x] + 1) \leq \phi([x]) + \phi(1) \leq [x] \phi(1) + \phi(1) \\ &\leq x \phi(1) + \phi(1) = (x + 1) \phi(1). \end{aligned}$$

If  $x > 1$ , then  $\phi(x) \leq 2x \phi(1)$ ; and if  $x \leq 1$ , then  $\phi(x) \leq \phi(1)$ .

Now, let  $f \in L^1(\mu, X)$ , and let  $A = \{ t \in T : \|f(t)\| \leq 1 \}$

$$B = \{ t \in T : \|f(t)\| > 1 \}$$

$$\begin{aligned} \|f\|_\phi &= \int_T \phi(\|f(t)\|) d\mu(t) \\ &= \int_A \phi(\|f(t)\|) d\mu(t) + \int_B \phi(\|f(t)\|) d\mu(t) \\ &\leq \int_T \phi(1) d\mu(t) + \int_T 2 \|f(t)\| \phi(1) d\mu(t) \\ &\leq \phi(1) \mu(T) + 2\phi(1) \|f\|_1 < \infty \quad \text{since } f \in L^1(\mu, X). \end{aligned}$$

Hence  $f \in L^\phi(\mu, X)$ . ■

## 2. Best Approximation in $L^\phi(\mu, X)$

In this section we investigate when  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  where  $G$  is a closed subspace of a Banach space  $X$ .

We present the following useful theorem.

**Theorem 3.2.1** : [4 , p.73] . Let  $G$  be a closed subspace of a Banach space  $X$  , if  $g$  is a best approximant of  $f$  in  $L^\phi(\mu, G)$  . Then  $g(t)$  is a best approximant of  $f(t)$  in  $G$  for almost all  $t \in T$  .

The following theorem establishes the relation between proximality in  $L^\phi(\mu, X)$  and  $L^1(\mu, X)$  .

**Theorem 3.2.2** : [4 , p.73] . Let  $G$  be a closed subspace of Banach space  $X$  , the following are equivalent :

- (i)  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .
- (ii)  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  .

**Proof** : (i)→(ii) . Let  $f \in L^1(\mu, X)$  , since  $L^1(\mu, X) \subset L^\phi(\mu, X)$  then  $f \in L^\phi(\mu, X)$  , but  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  so there exist  $g \in L^\phi(\mu, G)$  such that

$$\|f - g\|_\phi \leq \|f - h\|_\phi \quad \forall h \in L^\phi(\mu, G) .$$

Theorem (3.2.1) implies

$$\|f(t) - g(t)\| \leq \|f(t) - y\| \quad \forall y \in G \text{ for almost all } t \in T \dots (1)$$

Since  $0 \in G$  we get

$$\|f(t) - g(t)\| \leq \|f(t) - 0\| = \|f(t)\|$$

Hence

$$\begin{aligned} \|g(t)\| &= \|g(t) - f(t) + f(t)\| \\ &\leq \|g(t) - f(t)\| + \|f(t)\| \\ &\leq 2 \|f(t)\| \end{aligned}$$

Therefore  $g \in L^1(\mu, G)$ .

From (1) we get

$$\|f(t) - g(t)\| \leq \|f(t) - h(t)\| \quad \forall h \in L^\phi(\mu, G) \text{ a.e. } t$$

Integrating both sides we get

$$\|f - g\|_1 \leq \|f - h\|_1 \quad \forall h \in L^\phi(\mu, G).$$

Therefore  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$ .

Conversely, (ii)  $\rightarrow$  (i). Define the map  $J : L^\phi(\mu, X) \rightarrow L^1(\mu, X)$  by  $J(f) = \hat{f}$

where

$$\hat{f}(t) = \begin{cases} \frac{\phi(\|f(t)\|)}{\|f(t)\|} f(t) & f(t) \neq 0 \\ 0 & f(t) = 0 \end{cases}$$

At first we show that  $\hat{f} \in L^1(\mu, X)$

$$\|\hat{f}\|_1 = \int_T \|\hat{f}(t)\| d\mu(t)$$

$$\begin{aligned}
&= \int_{\mathcal{T}} \frac{\phi(\|f(t)\|)}{\|f(t)\|} \|f(t)\| \, d\mu(t) \\
&= \int_{\mathcal{T}} \phi(\|f(t)\|) \, d\mu(t) \\
&= \|f\|_{\phi} < \infty .
\end{aligned}$$

Second , we claim that  $J$  is onto .

$$\text{Let } g \in L^1(\mu, X) \text{ and let } f(t) = \begin{cases} \frac{\phi^{-1}(\|g(t)\|)}{\|g(t)\|} g(t) & g(t) \neq 0 \\ 0 & g(t) = 0 \end{cases}$$

$$\begin{aligned}
\text{Then } \|f\|_{\phi} &= \int_{\mathcal{T}} \phi(\|f(t)\|) \, d\mu(t) \\
&= \int_{\mathcal{T}} \phi\left[\frac{\phi^{-1}(\|g(t)\|)}{\|g(t)\|} \|g(t)\|\right] \, d\mu(t) \\
&= \int_{\mathcal{T}} \|g(t)\| \, d\mu(t) \\
&= \|g\|_1
\end{aligned}$$

Hence  $f \in L^{\phi}(\mu, X)$  and  $J(f) = g$  .

Finally since  $\phi$  is one-to-one it follows that  $J$  is one-to-one . It is now clear that

$$J(L^{\phi}(\mu, G)) = L^1(\mu, G) .$$

Now , let  $f \in L^{\phi}(\mu, X)$  . Then  $J(f) = \hat{f} \in L^1(\mu, X)$  and there exists  $\hat{g} \in L^1(\mu, G)$  such that  $\|\hat{f} - \hat{g}\|_1 \leq \|\hat{f} - \hat{h}\|_1$  for all  $\hat{h} \in L^1(\mu, G)$  . By Theorem (3.2.1) ; we have

$$\|\hat{f}(t) - \hat{g}(t)\| \leq \|\hat{f}(t) - y\| \text{ for all } y \in G \quad \text{a.e.t.}$$

Since  $\hat{g} \in L^1(\mu, G)$  and  $J$  is onto, there exists  $g \in L^\phi(\mu, G)$  s.t.  $J(g) = \hat{g}$ .

$$\text{Hence } \left\| f(t) - \frac{\phi(\|g(t)\|) \|f(t)\| g(t)}{\phi(\|f(t)\|) \|g(t)\|} \right\| \leq \left\| f(t) - \frac{\|f(t)\|}{\phi(\|f(t)\|)} y \right\| \quad \text{a.e.t}$$

and for all  $y \in G$ .

$$\text{Now take } h \in L^\phi(\mu, G). \text{ Then } \frac{\phi(\|f(t)\|)}{\|f(t)\|} h(t) \in G \quad \text{a.e.t}$$

Hence  $\|f(t) - w(t)\| \leq \|f(t) - h(t)\|$  a.e.t and for all  $h \in L^\phi(\mu, G)$ ,

$$\text{where } w(t) = \frac{\phi(\|g(t)\|) \|f(t)\| g(t)}{\phi(\|f(t)\|) \|g(t)\|}$$

using the fact that  $\|g(t)\| \leq 2 \|f(t)\|$  we will show that  $w \in L^\phi(\mu, G)$  as

follows

$$\begin{aligned} \|w(t)\| &= \frac{\phi(\|g(t)\|) \|f(t)\| \cdot \|g(t)\|}{\phi(\|f(t)\|) \|g(t)\|} \\ &\leq \frac{2\phi(\|f(t)\|) \cdot \|f(t)\|}{\phi(\|f(t)\|)} \\ &= 2 \|f(t)\| \end{aligned}$$

Hence  $w \in L^\phi(\mu, G)$ . Thus  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$ . ■

In a similar way we can prove the following theorem.

**Theorem 3.2.3** : [2, p.297]. Let  $G$  be a closed subspace of Banach space

$X$ . If  $1 < p < \infty$  the following are equivalent :

(i)  $L^p(\mu, G)$  is proximal in  $L^p(\mu, X)$ .



(ii)  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$ .

For  $p = \infty$ , we have :

**Theorem 3.2.4** : [1, p.528]. Let  $G$  be a closed subspace of Banach space  $X$ . If  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$ , then  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$ .

**Proof** : Let  $f \in L^\infty(\mu, X)$ . Since  $L^\infty(\mu, X) \subseteq L^1(\mu, X)$ , we have  $f \in L^1(\mu, X)$ .

By assumption, there exists  $f_1 \in L^1(\mu, G)$  such that

$$\|f - f_1\|_1 = d(f, L^1(\mu, G)).$$

By Theorem (3.2.1), it follows that

$$\|f(t) - f_1(t)\| = d(f(t), G) \text{ a.e.t.}$$

Hence  $\|f(t) - f_1(t)\| \leq \|f(t) - y\|$  a.e.t., and for all  $y \in G$ .

In particular

$$\|f(t) - f_1(t)\| \leq \|f(t) - g(t)\| \text{ a.e.t. , and for all } g \in L^1(\mu, G).$$

But  $L^\infty(\mu, G) \subseteq L^1(\mu, G)$ , and hence, for every  $h \in L^\infty(\mu, G)$  we have

$$\|f(t) - f_1(t)\| \leq \|f(t) - h(t)\| \text{ a.e.t. ....(*)}$$

Now, since  $0 \in G$ , we get  $\|f_1(t)\| \leq 2\|f(t)\|$  a.e.t. Hence  $f_1 \in L^\infty(\mu, G)$ .

Thus it follows from (\*) that

$$\|f - f_1\|_\infty \leq \|f - h\|_\infty \text{ for every } h \in L^\infty(\mu, G).$$

Consequently  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$ . ■

**Theorem 3.2.5** : Let  $G$  be a closed subspace of a Banach space  $X$ . If

$L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$ , then  $G$  is proximal in  $X$ .

**Proof :** Let  $x \in X$ , we define  $f(t) = x$  for all  $t \in T$ . Then  $f \in L^\phi(\mu, X)$ .

Since  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$ , there exists  $g \in L^\phi(\mu, G)$  such that

$\|f - g\|_\phi = d(f, L^\phi(\mu, G))$ . Theorem (3.2.1) implies  $\|f(t) - g(t)\| \leq \|f(t) - y\|$

a.e.t and for all  $y \in G$ . Hence  $\|x - g(t)\| \leq \|x - y\|$  for all  $y \in G$ .

Consequently  $G$  is proximal in  $X$ . ■

For  $p = \infty$ , we have :

**Theorem 3.2.6 :** Let  $G$  be a closed subspace of a Banach space  $X$ . If

$L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$ , then  $G$  is proximal in  $X$ .

**Proof :** Let  $x \in X$ . Consider the function  $f(t) = x$  for all  $t \in T$ . Then

$f \in L^\infty(\mu, X)$ . Hence there exists  $g \in L^\infty(\mu, G)$  s.t  $\|f - g\|_\infty = d(f, L^\infty(\mu, G))$ .

Theorem [11, p.36];  $\|f - g\|_\infty = \sup_t d(f(t), G)$ .

Hence  $\|f - g\|_\infty = \sup_t d(x, G)$ . Since  $f(t) = x$  for all  $t \in T$ .

$$\|f - g\|_\infty = d(x, G). \text{ But } d(x, G) = \sup \{ \|x - g(t)\| : t \in T \}.$$

$$\Rightarrow \|x - g(t)\| \leq d(x, G), \text{ for all } t \in T.$$

Therefore  $G$  is proximal in  $X$ . ■

For  $1 < p < \infty$ , we have :

**Theorem 3.2.7 :** [3, p.8]. Let  $G$  be a closed subspace of a Banach space

$X$ . If  $L^p(\mu, G)$  is proximal in  $L^p(\mu, X)$ , then  $G$  is proximal in  $X$  for

$1 < p < \infty$ .

**Proof :** If  $L^p(\mu, G)$  is proximal in  $L^p(\mu, X)$  . Theorem (3.2.3) implies  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  . Theorem (3.2.4) implies  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$  . Theorem (3.2.6) implies  $G$  is proximal in  $X$  . ■

**Definition 3.2.8 :** [1 , p.529] . A subspace  $G$  of a Banach space  $X$  is called 1-complemented in  $X$  if there is a closed subspace  $W$  in  $X$  such that  $X = G \oplus W$  and the projection  $P : X \rightarrow W$  is a contractive projection .

**Lemma 3.2.9 :** [1 , p.529] . If  $G$  is 1-complemented in  $X$  , then  $G$  is proximal in  $X$  .

**Proof :** Let  $X = G \oplus W$  and  $x \in X$  . Then  $x = g + w$  , where  $g \in G$  ,  $w \in W$  and  $\|w\| \leq \|x\|$  . We show that  $\|x - g\| \leq \|x - y\|$  for all  $y \in G$  . If possible assume that there exist  $g_1 \neq g \in G$  s.t  $\|x - g_1\| < \|x - g\|$  .

Set  $w_1 = x - g_1$  . By the uniqueness of the representation of  $x$  , we have  $w_1 \notin W$  .

Hence  $w_1 = g_2 + w_2$  , where  $g_2 \in G$  ,  $w_2 \in W$  and  $\|w_2\| \leq \|w_1\|$  . Therefore

$$x = w_1 + g_1 = (g_2 + w_2) + g_1 = (g_1 + g_2) + w_2 ,$$

and consequently  $g = g_1 + g_2$  and  $w = w_2$  . Thus

$$\|w\| = \|w_2\| \leq \|w_1\| .$$

But by assumption ,  $\|w_1\| = \|x - g_1\| < \|x - g\| = \|w\|$  .

This contradicts the assumption . Consequently  $\|x - g\| \leq \|x - y\|$  for all  $y \in G$  . Hence  $G$  is proximal in  $X$  . ■

**Remark 3.2.10** : If  $G$  is 1-complemented in  $X$  , then  $G$  may not be a Chebyshev subspace .

**Proof** : Let  $X = \mathbf{R}^2$  and  $G = \{ (g,g) : g \in \mathbf{R} \}$  with  $\|(x,y)\| = |x| + |y|$  .

Then  $G$  is not Chebyshev by Example (2.2.1) .

Now , let  $W = \{ (0,w) : w \in \mathbf{R} \}$  . Then

$$(x,y) = (x,x) + (0,y - x) .$$

Hence  $\mathbf{R}^2 = G \oplus W$  .

We define  $P : X \rightarrow W$  as

$$P(x,y) = P[ (x,x) + (0,y - x) ] = (0,y - x)$$

Now  $\|w\| = \|(0,y - x)\| = |y - x| \leq |y| + |x| = \|(x,y)\|$  .

Hence  $P$  is a contractive projection .

Therefore  $G$  is 1-complemented in  $\mathbf{R}^2$  .

**Theorem 3.2.11** : If  $G$  is 1-complemented in  $X$  , then  $L^1(\mu,G)$  is 1-complemented in  $L^1(\mu,X)$  .

**Proof** : Let  $X = G \oplus W$  and let  $P : X \rightarrow W$  be contractive projection . Hence

$x = (I - P)(x) + P(x)$  and  $\|P(x)\| \leq \|x\|$  . For  $f \in L^1(\mu,X)$  , set  $f_1 = (I - P)f$  ,

$f_2 = Pf$  a.e.t . Then

$$\|f_2\|_1 = \int_{\tau} \|f_2(t)\| d\mu(t) = \int_{\tau} \|P(f(t))\| d\mu(t) \leq \int_{\tau} \|f(t)\| d\mu(t) = \|f\|_1 < \infty .$$

Hence  $f_2 \in L^1(\mu,W)$  . Also

$$\begin{aligned}
\|f_1\|_1 &= \int_T \|f_1(t)\| \, d\mu(t) = \int_T \|(I-P)(f(t))\| \, d\mu(t) = \int_T \|f(t) - P(f(t))\| \, d\mu(t) \\
&\leq \int_T \|f(t)\| \, d\mu(t) + \int_T \|P(f(t))\| \, d\mu(t) \leq \int_T \|f(t)\| \, d\mu(t) + \int_T \|f(t)\| \, d\mu(t) \\
&= 2 \|f\|_1 < \infty .
\end{aligned}$$

Hence  $f_1 \in L^1(\mu, G)$  . Clearly  $f = f_1 + f_2$  .

Since  $W$  is a closed subspace of  $X$  , then  $L^1(\mu, W)$  is a closed subspace of  $L^1(\mu, X)$  . Also if  $f \in L^1(\mu, W) \cap L^1(\mu, G)$  then  $f \in L^1(\mu, W)$  and  $f \in L^1(\mu, G)$

$\Rightarrow f(t) \in W$  and  $f(t) \in G \quad \forall t \in T$  , but  $G \cap W = \{0\}$  .

$\Rightarrow f(t) = 0 \quad \forall t \in T \Rightarrow f = \hat{0}$  (zero function) .

Hence  $L^1(\mu, X) = L^1(\mu, G) \oplus L^1(\mu, W)$  . Define the map  $\hat{p} : L^1(\mu, X) \rightarrow L^1(\mu, W)$  by  $\hat{p}(f) = Pof = f_2$  for all  $f \in L^1(\mu, X)$  ,  $\hat{p}$  is a contractive projection . So  $L^1(\mu, G)$  is 1-complemented in  $L^1(\mu, X)$  . ■

**Corollary 3.2.12** : If  $G$  is 1-complemented in  $X$  , then  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  .

**Proof** : The corollary follows from the above Theorem and Lemma (3.2.9) . ■

**Definition 3.2.13** : [4 , p.72] . A closed subspace  $G$  of a Banach space  $X$  is called a  $\phi$ -summand of  $X$  if there is a bounded projection  $P : X \rightarrow G$  such that

$$\phi(\|x\|) = \phi(\|P(x)\|) + \phi(\|x - P(x)\|) \text{ for all } x \in X .$$

where  $\phi$  is a modulus function .

**Theorem 3.2.14 :** [4 , p.72] . If  $G$  is a  $\phi$ -summand of Banach space  $X$  , then  $G$  is proximal in  $X$  .

**Proof :** Let  $x \in X$  , for every  $g \in G$  we have

$$\begin{aligned}\phi (\|x - g\|) &= \phi (\|P(x-g)\|) + \phi (\|(x - g) - P(x-g)\|) \\ &= \phi (\|P(x) - g\|) + \phi (\|x - P(x)\|) \\ &\geq \phi (\|x - P(x)\|)\end{aligned}$$

Since  $\phi^{-1}$  exists and strictly increasing . Hence  $\|x - g\| \geq \|x - P(x)\|$  .

i.e  $P(x) \in P(x,G)$  . Thus  $G$  is proximal in  $X$  . ■

**Remark 3.2.15 :** If  $G$  is a  $\phi$ -summand of Banach space  $X$  , then  $G$  is a Chebyshev subspace .

**Proof :** Assume that  $G$  is a  $\phi$ -summand of  $X$  . Theorem (3.2.14) implies  $P(x) \in P(x,G)$  .

Now suppose  $g^*$  is another closest element to  $x$  .

i.e  $\|x - g^*\| = \|x - P(x)\|$  .....\*

But  $x - g^* \in X$

$$\begin{aligned}\phi (\|x - g^*\|) &= \phi (\|P(x - g^*)\|) + \phi (\|x - g^* - P(x - g^*)\|) \\ &= \phi (\|P(x) - g^*\|) + \phi (\|x - P(x)\|)\end{aligned}$$

by (\*)  $\phi (\|P(x) - g^*\|) = 0$  . So ,  $P(x) = g^*$  .

Therefore  $P(x)$  is the unique closest element .

Thus  $G$  is a Chebyshev subspace . ■

**Theorem 3.2.16** : [4 , p.73] . Let  $G$  be a proximal subspace of Banach space  $X$  . Then for every simple function  $f \in L^\phi(\mu, X)$  ,  $P(f, L^\phi(\mu, G))$  is not empty .

**Proof** : Let  $f = \sum_{i=1}^n a_i \chi_{E_i}$  where  $E_i$  are disjoint measurable sets in  $T$  .

Set  $g = \sum_{i=1}^n b_i \chi_{E_i}$  where  $b_i \in P(a_i, G)$  . If  $h$  is an element in  $L^\phi(\mu, G)$  , then

$$\begin{aligned}
 \|f - h\|_\phi &= \int_T \phi(\|f(t) - h(t)\|) d\mu(t) \\
 &= \sum_{i=1}^n \int_{E_i} \phi(\|f(t) - h(t)\|) d\mu(t) \\
 &= \sum_{i=1}^n \int_{E_i} \phi(\|a_i - h(t)\|) d\mu(t) \\
 &\geq \sum_{i=1}^n \int_{E_i} \phi(\|a_i - b_i\|) d\mu(t) \\
 &= \int_T \phi(\|f(t) - g(t)\|) d\mu(t) \\
 &= \|f - g\|_\phi
 \end{aligned}$$

Therefore  $\|f - g\|_\phi \leq \|f - h\|_\phi \quad \forall h \in L^\phi(\mu, G)$  . Thus  $g \in P(f, L^\phi(\mu, G))$  . ■

## Chapter 4

### S-property

**Introduction :** Let  $X$  be a Banach space and  $G$  a closed subspace of  $X$ . The space  $G$  is said to have the S-property if  $z_1 \in P(x_1, G)$  and  $z_2 \in P(x_2, G)$  then  $z_1 + z_2 \in P(x_1 + x_2, G)$  for all  $x_1, x_2 \in X$ . In this chapter we prove that if  $G$  has the S-property, then  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  if and only if  $G$  is proximal in  $X$ . As an application of this result we prove that  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  if and only if  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$ , in case  $G$  has the S-property.

An example of a subspace  $G \subset X$  which has the S-property is the following.

**Example 4.1.1 :** Let  $X = \mathbb{R}^2$ . Set  $G = \{ (x, 0) : x \in \mathbb{R} \}$  with the Euclidean norm.

Now if  $x_1 = (m, n)$ , then  $P(x_1, G) = \{(m, 0)\}$  i.e.  $z_1 = (m, 0)$ .

and if  $x_2 = (r, s)$ , then  $P(x_2, G) = \{(r, 0)\}$  i.e.  $z_2 = (r, 0)$ .

But  $x_1 + x_2 = (m+r, n+s)$ ; so  $P(x_1 + x_2, G) = \{(m+r, 0)\}$  i.e.

$$(m+r, 0) = z_1 + z_2.$$

Therefore  $G$  has the S-property.

**Remark 4.1.2 :** In a Banach space  $X$ , if  $G$  has the S-property, it does not necessarily follow that  $G$  is proximal in  $X$ . For example, if  $X = c_0$  with



$\|x\| = \sup_n |x_n|$  and  $G = \{ x = (x_n) \in c_0 : \sum_{n=1}^{\infty} 2^{-n} x_n = 0 \}$ . Remark (2.3.1)

implies  $G$  is not proximal in  $c_0$ .

Now let  $x \in c_0 - G$  and suppose that  $P(x, G) \neq \emptyset$  i.e.  $\exists z \in G$  s.t.  $z \in P(x, G)$ ;

so  $0 \in P(x - z, G)$  by part (2) of Theorem (2.2.5). This means there exists

$x - z \in c_0 - \{0\}$  such that  $0 \in P(x - z, G)$ . Theorem (2.3.7) implies  $G$  is

proximal in  $c_0$  which is a contradiction. Therefore  $P(x, G) = \emptyset$  for every

$$x \in c_0 - G.$$

Hence  $G$  has the S-property.

We shall now give various closed subspaces of a Banach space which have the S-property.

**Theorem 4.1.3 :** If  $G$  is a  $\phi$ -summand of  $X$ , then  $G$  has the S-property.

**Proof :** Let  $z_1 \in P(x_1, G)$ ,  $z_2 \in P(x_2, G)$ . Since  $G$  is a  $\phi$ -summand of  $X$ , then there exist a projection  $E : X \rightarrow G$  s.t.  $E(x)$  is a unique best approximant of  $x$  in  $G$  for all  $x \in X$  by Theorem (3.2.14) and Remark (3.2.15). Hence

$$z_1 = E(x_1) \text{ and } z_2 = E(x_2).$$

But  $z_1 + z_2 = E(x_1) + E(x_2) = E(x_1 + x_2)$  since  $E$  is linear.

This implies that  $z_1 + z_2 \in P(x_1 + x_2, G)$ . Thus  $G$  has the S-property. ■

**Theorem 4.1.4 :** If  $G$  is 1-complemented and Chebyshev in  $X$ , then  $G$  has the S-property.

**Proof :** Let  $z_1 \in P(x_1, G)$ ,  $z_2 \in P(x_2, G)$ , since  $G$  is 1-complemented in  $X$ , there is a closed subspace  $W \subset X$  s.t  $X = G \oplus W$ . This implies  $x_1$  and  $x_2$  can be written, uniquely, in the form

$$x_1 = g_1 + w_1, \quad x_2 = g_2 + w_2$$

where  $g_1, g_2 \in G$ , and  $w_1, w_2 \in W$ .

From the proof of Lemma (3.2.9) and the assumption that  $G$  is Chebyshev, we get that  $z_1 = g_1, z_2 = g_2$ .

Now  $x_1 + x_2 = (g_1 + g_2) + (w_1 + w_2)$ .

Since  $G$  is a subspace, so  $g_1 + g_2 \in G$ .

Also  $W$  is a subspace, so  $w_1 + w_2 \in W$ .

It now follows that  $z_1 + z_2 = g_1 + g_2 \in P(x_1 + x_2, G)$ .

Thus  $G$  has the S-property. ■

**Theorem 4.1.5 :** Let  $G$  be a closed subspace of a Hilbert space  $X$ , then  $G$  has the S-property.

**Proof :** Let  $z_1 \in P(x_1, G)$  and  $z_2 \in P(x_2, G)$ . We show  $z_1 + z_2 \in P(x_1 + x_2, G)$ .

Theorem (2.2.6) implies  $x_1 - z_1 \perp G$ , and  $x_2 - z_2 \perp G$ . Hence

$$\langle x_1 - z_1, g \rangle = 0 \text{ and } \langle x_2 - z_2, g \rangle = 0 \text{ for all } g \in G.$$

Now  $\langle x_1 + x_2 - (z_1 + z_2), g \rangle = \langle x_1 - z_1, g \rangle + \langle x_2 - z_2, g \rangle = 0$  for all  $g \in G$ .

Hence  $x_1 + x_2 - (z_1 + z_2) \perp G$ . Theorem (2.2.6) implies

$$z_1 + z_2 \in P(x_1 + x_2, G)$$

Thus  $G$  has the  $S$ -property . □

**Theorem 4.1.6 :** If  $G$  is a semi-Chebyshev hyperplane in a Banach space  $X$  passing through zero , then  $G$  has the  $S$ -property .

**Proof :** Case (1) : If  $G$  is proximal in  $X$  . Let  $f \in X^*$  so that

$G = \{ y \in X , f(y) = 0 \}$  . Fix an arbitrary  $z \in X \setminus G$  so ;  $f(z) \neq 0$  , and let

$$y_0 = x - f(x)/f(z) \cdot z \quad \text{where } x \in X$$

So  $f(y_0) = 0$  , whence  $y_0 \in G$  .Consequently

$$X = G \oplus W \quad \text{where } W = \{ w = \alpha z : \alpha \text{ scalar} \} \dots\dots\dots*$$

Now let  $z_1 \in P(x_1, G)$  ,  $z_2 \in P(x_2, G)$  . It will be shown that

$$z_1 + z_2 \in P(x_1 + x_2, G) .$$

By (\*) every  $x_1 \in X$  ,  $x_2 \in X$  can be written , uniquely , in the form

$$x_1 = g_1 + \alpha_1 z , \quad x_2 = g_2 + \alpha_2 z \dots\dots\dots**$$

where  $g_1 , g_2 \in G$  and  $\alpha_1 , \alpha_2$  are scalars .

Now assume that  $g' \in P(x_1 + x_2, G)$  , then by \*\*

$g' \in P(g_1 + g_2 + (\alpha_1 + \alpha_2)z, G)$  . Theorem (2.2.5) implies

$$\begin{aligned} g' &= (g_1 + g_2) + (\alpha_1 + \alpha_2) w \quad \text{where } w \in P(z, G) \\ &= g_1 + \alpha_1 w + g_2 + \alpha_2 w \end{aligned}$$

Since  $w \in P(z, G)$  , Theorem (2.2.5) implies

$$g_1 + \alpha_1 w \in P(g_1 + \alpha_1 z, G) = P(x_1, G) .$$

And  $g_2 + \alpha_2 w \in P(g_2 + \alpha_2 z, G) = P(x_2, G)$ .

Hence

$$g_1 + \alpha_1 w = z_1$$

$$g_2 + \alpha_2 w = z_2$$

Consequently  $g' = z_1 + z_2$ .

Therefore  $z_1 + z_2 \in P(x_1 + x_2, G)$ . Thus  $G$  has the S-property.

Case (2) : If  $G$  is not proximal in  $X$ . Theorem (2.3.7) implies  $P(x, G) = \phi$  for every  $x \in X - G$ . Thus  $G$  has the S-property. ■

We now state and prove an important result for  $G$  with S-property.

**Theorem 4.1.7** : Let  $X$  be any Banach space, and  $G$  a closed subspace of  $X$  which has the S-property, then  $P_G^{-1}(0)$  is a closed subspace of  $X$  and

$$P_G^{-1}(0) \cap G = \{0\}.$$

**Proof** : Let  $x_1, x_2 \in P_G^{-1}(0)$ , so  $0 \in P(x_1, G)$  and  $0 \in P(x_2, G)$ . Since  $G$  has the S-property we get  $0 \in P(x_1 + x_2, G)$ . Hence  $x_1 + x_2 \in P_G^{-1}(0)$ . \*

Let  $x \in P_G^{-1}(0)$  and  $\alpha$  be any scalar. Then

$$d(\alpha x, G) = |\alpha| d(x, G) = |\alpha| \|x\| = \|\alpha x\| \Rightarrow 0 \in P(\alpha x, G) \Rightarrow \alpha x \in P_G^{-1}(0). **$$

By (\*), (\*\*),  $P_G^{-1}(0)$  is a subspace of  $X$ .

Now let  $(x_n)$  be a sequence in  $P_G^{-1}(0)$  and  $x \in X$  such that  $\lim x_n = x$ . Since

$G$  is a subspace of  $X$ ,  $0 \in G$ ,  $d(x, G) \leq \|x\| \dots\dots(1)$ .

Given  $\epsilon > 0$  there exist a natural number  $N(\epsilon)$  such that  $\|x_n - x\| < \epsilon$  for all  $n > N(\epsilon)$ .

Fix  $n \geq N(\epsilon)$  to have :

$$\begin{aligned}
 \|x\| &= \|x - x_n + x_n\| \leq \|x - x_n\| + \|x_n\| \\
 &< \epsilon + | \|x_n\| - d(x,G) + d(x,G) | \\
 &\leq \epsilon + | \|x_n\| - d(x,G) | + d(x,G) \\
 &= \epsilon + | d(x_n,G) - d(x,G) | + d(x,G) \quad \text{since } x_n \in P_G^{-1}(0) \\
 &\leq \epsilon + \|x_n - x\| + d(x,G) \\
 &\leq 2\epsilon + d(x,G)
 \end{aligned}$$

Hence  $\|x\| \leq d(x,G) \dots\dots(2)$

From (1) , (2) we get  $\|x\| = d(x,G)$  . So ,  $0 \in P(x,G)$  hence  $x \in P_G^{-1}(0)$  .

Thus  $P_G^{-1}(0)$  is closed .

Let  $g \in P_G^{-1}(0) \cap G \Rightarrow g \in P_G^{-1}(0)$  and  $g \in G$  .

$$\Rightarrow 0 \in P(g,G) \text{ and } g \in G .$$

$$\Rightarrow \|g\| = d(g,G) \text{ and } g \in G .$$

$$\Rightarrow \|g\| = 0 .$$

$$\Rightarrow g = 0$$

Therefore  $P_G^{-1}(0) \cap G = \{0\}$  . ■

The next theorem shows that if we add the condition that  $G$  has the S-property then the converse of Lemma (3.2.9) will be true .

**Theorem 4.1.8** : Let  $X$  be any Banach space , and  $G$  a closed subspace of  $X$  which has the S-property .  $G$  is proximal in  $X$  if and only if  $G$  is 1-complemented in  $X$  .

**Proof** : If  $G$  is 1-complemented in  $X$  , then by Lemma (3.2.9) it is proximal in  $X$  .

Suppose now that  $G$  is proximal in  $X$  . Theorem (2.3.2) implies

$X = G + P_G^{-1}(0)$  . Theorem (4.1.7) shows that  $P_G^{-1}(0)$  is a closed subspace of  $X$  and  $P_G^{-1}(0) \cap G = \{0\}$  . Hence  $X = G \oplus P_G^{-1}(0)$  .

Now define  $P : X \rightarrow P_G^{-1}(0)$  by

$$P(x) = P(g + z) = z \text{ where } x = g + z, g \in G, z \in P_G^{-1}(0) .$$

$$\begin{aligned} \|x\| &\geq d(x, G) = d(g + z, G) \\ &= d(z, G) \\ &= \|z\| \end{aligned}$$

Therefore  $\|x\| \geq \|z\|$  .

Hence  $P$  is a contractive projection . Thus  $G$  is 1-complemented in  $X$  . ■

An important application of the previous theorem is the following .

**Theorem 4.1.9** : Let  $X$  be any Banach space , and  $G$  be a closed subspace of  $X$  which has the S-property . The following are equivalent :

- (i)  $G$  is proximal in  $X$  .
- (ii)  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  .

(iii)  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .

**Proof :** (i)→(ii) , assume  $G$  is proximal in  $X$  . Theorem (4.1.8) implies  $G$  is 1-complemented in  $X$  . Corollary (3.2.12) implies  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  .

(ii)→(iii) , assume  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  . Theorem (3.2.2) implies  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .

(iii)→(i) , assume that  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$ . Theorem (3.2.5) implies  $G$  is proximal in  $X$  . ■

We shall now give various corollaries of this theorem .

**Corollary 4.1.10 :** Let  $X$  be a Banach space , and  $G$  is a  $\phi$ -summand of  $X$  , then  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .

**Proof :** The corollary follows from the above Theorem and Theorems (4.1.3) , (3.2.14) . ■

**Corollary 4.1.11 :** Let  $G$  be a closed subspace of a Hilbert space  $X$  , then  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .

**Proof :** The corollary follows from the above Theorem and Theorems (2.4.3) , (4.1.5) . ■

**Corollary 4.1.12 :** If  $G$  is a Chebyshev hyperplane in a Banach space  $X$  passing through zero , then  $L^\phi(\mu, G)$  is proximal in  $L^\phi(\mu, X)$  .

**Proof :** The corollary follows from the above Theorem and Theorem (4.1.6) . ■

In particular , it follows that every Orlicz space  $L^\phi(\mu, \bar{X})$  where  $X$  is a Banach space and strictly convex has at least one proximal linear subspace .

Now we state and prove our main result .

**Theorem 4.1.13 :** Let  $X$  be any Banach space , and  $G$  be a closed subspace of  $X$  with the S-property . Then  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  if and only if  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$  .

**Proof :** If  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  , then by Theorem (3.2.4)  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$  .

Conversely if  $L^\infty(\mu, G)$  is proximal in  $L^\infty(\mu, X)$  , then  $G$  is proximal in  $X$  by Theorem (3.2.7) .

Theorem (4.1.9) , implies  $L^1(\mu, G)$  is proximal in  $L^1(\mu, X)$  . ■

## 2. Further Results :-

**Theorem 4.2.1 :** Let  $X$  be a Banach space and  $G$  be a closed subspace of  $X$  , if  $G$  has the S-property in  $X$  , then  $L^\phi(\mu, G)$  has the S-property in  $L^\phi(\mu, X)$  .

**Proof :** Let  $g_1 \in P(f_1, L^\phi(\mu, G))$  and  $g_2 \in P(f_2, L^\phi(\mu, G))$  , we will show that :

$$g_1 + g_2 \in P(f_1 + f_2 , L^\phi(\mu, G))$$

Now

$$g_1 \in P(f_1, L^\phi(\mu, G)) . \text{ Theorem (3.2.1) implies}$$



$$g_1(t) \in P(f_1(t), G) \quad \text{for all almost } t \in T \dots\dots(1)$$

Also

$$g_2 \in P(f_2, L^\phi(\mu, G)) . \text{ Theorem (3.2.1) implies}$$

$$g_2(t) \in P(f_2(t), G) \quad \text{for all almost } t \in T \dots\dots\dots(2)$$

Since  $G$  has the S-property , from (1) and (2) we get

$$(g_1 + g_2)(t) \in P((f_1 + f_2)(t), G) \quad \text{for all almost } t \in T .$$

Hence

$$d((f_1 + f_2)(t), G) = \|(f_1 + f_2)(t) - (g_1 + g_2)(t)\|$$

$$\text{Hence } \|(f_1 + f_2)(t) - (g_1 + g_2)(t)\| \leq \|(f_1 + f_2)(t) - y\| \quad \text{a.e.t, and } \forall y \in G$$

In particular

$$\|(f_1 + f_2)(t) - (g_1 + g_2)(t)\| \leq \|(f_1 + f_2)(t) - h(t)\| \quad \text{a.e.t, and } \forall h \in L^\phi(\mu, G)$$

Since  $\phi$  is strictly increasing , then

$$\phi(\|(f_1 + f_2)(t) - (g_1 + g_2)(t)\|) \leq \phi(\|(f_1 + f_2)(t) - h(t)\|) \quad \text{a.e.t and}$$

$$\forall h \in L^\phi(\mu, G)$$

Integrating both sides we get

$$\|(f_1 + f_2) - (g_1 + g_2)\|_\phi \leq \|(f_1 + f_2) - h\|_\phi \quad \forall h \in L^\phi(\mu, G)$$

$$\text{Hence } d(f_1 + f_2, L^\phi(\mu, G)) = \|(f_1 + f_2) - (g_1 + g_2)\|_\phi$$

$$\text{Therefore } g_1 + g_2 \in P(f_1 + f_2, L^\phi(\mu, G)) .$$

Thus  $L^\phi(\mu, G)$  has the S-property . ■

The following theorem describes the relationship between  $P_{L^\phi(\mu, G)}^{-1}(0)$  and

$L^\phi(\mu, P_G^{-1}(0))$  .

**Theorem 4.2.2 :** Let  $X$  be a Banach space , and  $G$  be a closed subspace of  $X$  . If  $G$  has the S-property , then  $P_{L^\phi(\mu, G)}^{-1}(0) = L^\phi(\mu, P_G^{-1}(0))$  .

**Proof :** Let  $f \in L^\phi(\mu, P_G^{-1}(0))$  . This means  $f(t) \in P_G^{-1}(0)$  , and that  $\|f\|_\phi < \infty$  .

Now

$$f(t) \in P_G^{-1}(0) ; \text{ so , } 0 \in P(f(t), G) ; \text{ hence } d(f(t), G) = \|f(t)\| .$$

$$\text{i.e } \|f(t)\| \leq \|f(t) - g\| \quad \forall g \in G$$

In particular

$$\|f(t)\| \leq \|f(t) - h(t)\| \quad \forall h \in L^\phi(\mu, G)$$

since  $\phi$  is strictly increasing , then

$$\phi(\|f(t)\|) \leq \phi(\|f(t) - h(t)\|) \quad \forall h \in L^\phi(\mu, G)$$

Integrating both sides we get

$$\|f\|_\phi \leq \|f - h\|_\phi \quad \forall h \in L^\phi(\mu, G)$$

$$\text{Hence } d(f, L^\phi(\mu, G)) = \|f\|_\phi$$

$$\text{Therefore } 0 \in P(f, L^\phi(\mu, G)) \Rightarrow f \in P_{L^\phi(\mu, G)}^{-1}(0)$$

Thus

$$L^\phi(\mu, P_G^{-1}(0)) \subset P_{L^\phi(\mu, G)}^{-1}(0) \quad \dots\dots\dots(1)$$

Let  $f \in P_{L^\phi(\mu, G)}^{-1}(0)$  . We claim that  $f \in L^\phi(\mu, P_G^{-1}(0))$  i.e.  $\|f\|_\phi < \infty$  and

$$f(t) \in P_G^{-1}(0) .$$

Now

$$f \in P_{L^\phi(\mu, G)}^{-1}(0) \subset L^\phi(\mu, X) . \text{ Hence } \|f\|_\phi < \infty \dots\dots\dots(2)$$

Also  $0 \in P(f, L^\phi(\mu, G))$  . Theorem (3.2.1) implies

$$0 \in P(f(t), G) \text{ a.e.t.}$$

$$\text{Thus } f(t) \in P_G^{-1}(0) \dots\dots\dots(3)$$

From (2) and (3) we get

$$f \in L^\phi(\mu, P_G^{-1}(0))$$

Therefore

$$P_{L^\phi(\mu, G)}^{-1}(0) \subset L^\phi(\mu, P_G^{-1}(0)) \dots\dots\dots(4)$$

From (1) and (4) we get  $P_{L^\phi(\mu, G)}^{-1}(0) = L^\phi(\mu, P_G^{-1}(0))$  . ■

**Lemma 4.2.3 :** Let  $X$  be a Banach space , and  $G_1 , G_2$  are closed subspaces of  $X$  . If  $G_1 \subset G_2$  then  $P_{G_2}^{-1}(0) \subset P_{G_1}^{-1}(0)$

**Proof :** Let  $x \in P_{G_2}^{-1}(0) \Rightarrow 0 \in P(x, G_2) \Rightarrow d(x, G_2) = \|x\|$  , but  $G_1 \subset G_2 \Rightarrow \|x\| \geq d(x, G_1) \geq d(x, G_2) = \|x\|$  .

Therefore  $d(x, G_1) = \|x\|$  . Hence  $x \in P_{G_1}^{-1}(0)$  . ■

**Theorem 4.2.4 :** Let  $X$  be a Banach space , and  $G$  a closed subspace of  $X$  which has the S-property . If  $G$  is proximal in  $X$  then  $P_G^{-1}(0)$  is proximal in  $X$  and has the S-property .

**Proof :** Let  $x \in X$ . The proof of Theorem (4.1.8) implies  $x$  can be written , uniquely , in the form

$$x = g + z \text{ where } g \in G \text{ and } z \in P_G^{-1}(0) \dots\dots\dots(1)$$

$$\text{Now } g \in G \Rightarrow g \perp w \quad \forall w \in P_G^{-1}(0) \Rightarrow g \perp P_G^{-1}(0) \Rightarrow 0 \in P(g, P_G^{-1}(0))$$

$$\Rightarrow d(g, P_G^{-1}(0)) = \|g\| \dots\dots\dots(2)$$

From (1) and (2) we get

$$d(x - z, P_G^{-1}(0)) = \|x - z\| \Rightarrow d(x, P_G^{-1}(0)) = \|x - z\|$$

$$\text{Therefore } z \in P(x, P_G^{-1}(0)) \text{ i.e } z = x - g \text{ where } g \in P(x, G) \dots\dots\dots(3)$$

Thus  $P_G^{-1}(0)$  is proximal in  $X$ .

To show  $P_G^{-1}(0)$  has the S-property .

$$\text{Let } z_1 \in P(x_1, P_G^{-1}(0)) \text{ and } z_2 \in P(x_2, P_G^{-1}(0))$$

$$\text{From (3) we get } x_1 - z_1 \in P(x_1, G) \text{ and } x_2 - z_2 \in P(x_2, G)$$

Since  $G$  has the S-property , then

$$x_1 + x_2 - (z_1 + z_2) \in P(x_1 + x_2, G)$$

$$\Rightarrow z_1 + z_2 \in P(x_1 + x_2, P_G^{-1}(0))$$

Thus  $P_G^{-1}(0)$  has the S-property . ■

**Theorem 4.2.5 :** Let  $X$  be a Banach space , and  $G$  is proximal in  $X$ . If  $G$

has the S-property then  $P_{P_G^{-1}(0)}^{-1}(0) = G$  .

**Proof :** Let  $g \in G \Rightarrow z \perp g \quad \forall z \in P_G^{-1}(0) \Rightarrow g \perp P_G^{-1}(0)$

$\Rightarrow 0 \in P(g, P_G^{-1}(0)) \Rightarrow g \in P_{P_G^{-1}(0)}^{-1}(0)$  .

Therefore  $G \subset P_{P_G^{-1}(0)}^{-1}(0)$  .....(1)

Now , let  $x \in P_{P_G^{-1}(0)}^{-1}(0)$  . Then by the proof of Theorem (4.1.8) we have

$$x = x_1 + x_2 \quad \text{where } x_1 \in G \quad \text{and } x_2 \in P_G^{-1}(0) .$$

Since  $G \subset P_{P_G^{-1}(0)}^{-1}(0)$  ,  $x_1 \in P_{P_G^{-1}(0)}^{-1}(0)$  . Then  $x_2 = x - x_1 \in P_{P_G^{-1}(0)}^{-1}(0)$  .

But  $x_2 \in P_G^{-1}(0)$  . Theorem (4.1.8) implies  $x_2 = x - x_1 = 0$  .

$$\Rightarrow x = x_1 \Rightarrow x \in G$$

Therefore  $P_{P_G^{-1}(0)}^{-1}(0) \subset G$  .....(2)

Thus we proved that  $G = P_{P_G^{-1}(0)}^{-1}(0)$  . ■

**Theorem 4.2.6 :** Let  $X$  be a Banach space , and  $G$  is proximal in  $X$  . If  $G$  has the S-property then  $L^\phi(\mu, G) = L^1(\mu, G)$  .

**Proof :** Theorem (3.1.4) implies  $L^1(\mu, G) \subset L^\phi(\mu, G)$  .....(1)

also  $L^1(\mu, P_G^{-1}(0)) \subset L^\phi(\mu, P_G^{-1}(0))$  .

Lemma (4.2.3) implies  $P_{L^\phi(\mu, P_G^{-1}(0))}^{-1}(0) \subset P_{L^1(\mu, P_G^{-1}(0))}^{-1}(0)$  .

Theorem (4.2.2) implies  $L^\phi(\mu, P_{P_G^{-1}(0)}^{-1}(0)) \subset L^1(\mu, P_{P_G^{-1}(0)}^{-1}(0))$  .

Theorem (4.2.5) implies  $L^\phi(\mu, G) \subset L^1(\mu, G)$  .....(2)

From (1) and (2) we get  $L^\phi(\mu, G) = L^1(\mu, G)$ . ■

Let  $X$  be a Banach space, and  $G$  be proximal in  $X$ , then any map which associates with each element of  $X$  one of its best approximant in  $G$  is called a proximity map. This mapping is, in general, non linear.

**Theorem 4.2.7 :** Let  $X$  be a Banach space, and  $G$  be a Chebyshev subspace of  $X$ . There exists a linear proximity map if and only if  $G$  has the S-property.

**Proof :** Let  $E$  be a linear proximity map. we claim that  $G$  has the S-property.

Let  $z_1 \in P(x_1, G)$ , and  $z_2 \in P(x_2, G)$ . We show that

$$z_1 + z_2 \in P(x_1 + x_2, G)$$

Now

$$z_1 + z_2 = E(x_1) + E(x_2) = E(x_1 + x_2) \in P(x_1 + x_2, G).$$

Therefore  $G$  has the S-property.

Conversely, assume that  $G$  has the S-property.

Define  $E : X \rightarrow G$  such that  $E(x) \in P(x, G)$ .

Now, we claim that  $E$  is linear.

Let  $x_1, x_2 \in X$ , we show that  $E(x_1 + x_2) = E(x_1) + E(x_2)$ .

Now,  $E(x_1) \in P(x_1, G)$  and  $E(x_2) \in P(x_2, G)$ .

Since  $G$  has the S-property, then  $E(x_1) + E(x_2) \in P(x_1 + x_2, G)$ .

Also  $E(x_1 + x_2) \in P(x_1 + x_2, G)$ . Since  $G$  is a Chebyshev subspace then

$$E(x_1 + x_2) = E(x_1) + E(x_2) \dots\dots\dots(1)$$

Let  $x \in X$ ,  $\alpha$  scalar then  $E(x) \in P(x, G)$ . Theorem (2.2.5) implies

$$\alpha \cdot E(x) \in P(\alpha x, G), \text{ also } E(\alpha x) \in P(\alpha x, G).$$

Since  $G$  is a Chebyshev subspace of  $X$  then

$$E(\alpha x) = \alpha \cdot E(x) \dots\dots\dots(2)$$

By (1) and (2)  $E$  is a linear. ■

If  $S$  is a compact Hausdorff space and  $X$  is a Banach space, then  $C(S, X)$  will denote the Banach space of all continuous maps  $f$  from  $S$  into  $X$  with norm defined as  $\|f\| = \sup_s \|f(s)\|$ .

**Theorem 4.2.8** : Let  $S$  be a compact Hausdorff space, and  $G$  be a Chebyshev subspace of a Banach space  $X$  which has the  $S$ -property. If  $C(S, G)$  be a Chebyshev subspace of  $C(S, X)$  then  $C(S, G)$  has the  $S$ -property in  $C(S, X)$ .

**Proof** : Since  $G$  is a Chebyshev and has the  $S$ -property. Theorem (4.2.7) implies there exists a linear proximity map  $E : X \rightarrow G$ . Then define the map.

$$F : C(S, X) \rightarrow C(S, G) \text{ by}$$

$$F(f) = E \circ f.$$

Our claim is that  $F$  is a linear proximity map.

If  $g \in C(S,G)$  then

$$\|f(s) - E(f(s))\| \leq \|f(s) - g(s)\|$$

For all  $s \in S$ ,  $f \in C(S,X)$ . Hence

$$\|f - E \circ f\| \leq \|f - g\| \text{ for all } g \in C(S,G)$$

Therefore

$$F(f) \in P(f, C(S,G))$$

And consequently

$F$  is a proximity map .

Let  $f, g \in C(S,X)$ , we claim that  $F(f + g) = F(f) + F(g)$

$$\begin{aligned} F(f + g)(s) &= E(f(s) + g(s)) \\ &= E(f(s)) + E(g(s)) && \text{since } E \text{ is linear} \\ &= E \circ f(s) + E \circ g(s) \\ &= (E \circ f + E \circ g)(s) \end{aligned}$$

Hence  $F(f + g) = E \circ f + E \circ g$

$$= F(f) + F(g) \dots\dots\dots(1)$$

Let  $f \in C(S,X)$ , and  $\alpha$  (scalar)

$$\begin{aligned} F(\alpha f)(t) &= E(\alpha f(t)) \\ &= \alpha E(f(t)) && \text{since } E \text{ is linear} \\ &= \alpha [(E \circ f)(t)] \\ &= \alpha (F(f))(t) \end{aligned}$$



$$F(\alpha f) = \alpha F(f) \dots\dots\dots(2)$$

By (1) and (2)  $F$  is linear .

Theorem (4.2.7 ) implies  $C(S,G)$  has the  $S$ -property . ■

In [7 , p.333] , it is shown that if  $X$  is a Hilbert space , then  $X$  is strictly convex . In the following , we give another proof .

**Theorem 4.2.9 :** If  $X$  is a Hilbert space , then  $X$  is strictly convex .

**Proof :** Assume that  $X$  is not strictly convex , whence there exists an  $f \in X^*$  with  $\|f\| = 1$  which has two distinct maximal elements  $x,y$  of norm  $\|x\| = \|y\| = 1$  by Definition (1.3.3) .

Put  $G = [x - y] = \{ \alpha(x - y) : \alpha \text{ scalar} \}$ .

$$\|x\| = |f(x)| = |f(x - g)| \leq \|f\| \|x - g\| = \|x - g\| \quad \text{for all } g \in G .$$

Hence  $0 \in P(x,G)$  .

$$\text{Also } \|y\| = |f(y)| = |f(y - g)| \leq \|f\| \|y - g\| = \|y - g\| \quad \text{for all } g \in G .$$

Thus  $0 \in P(y,G) \Rightarrow 0 \in P(-y,G)$  .

But  $G$  is a closed subspace of a Hilbert space . Theorem (4.1.5) implies  $G$  has the  $S$ -property . Then  $0 \in P(x - y,G)$  which contradicts the fact  $x - y \in G - \{0\}$  . Therefore  $X$  is strictly convex . ■

**References :-**

1. Deeb , W. and Khalil , R. Best approximation in  $L(X,Y)$  , Math. Proc. Cambridge Philos . Soc. 104 (1988) , 527-531 .
2. Deeb , W. and Khalil , R. Best approximation in  $L^p(\mu,X)$  II , J. Approx. Theory , 59 (1989) , 296-299 .
3. Deeb , W. and Khalil , R. Best approximation in  $L^p(I,X)$  , III , 1994 .
4. Deeb , W. and Khalil , R. Best approximation in  $L^p(I,X)$  ,  $0 < p < 1$  , J. Approx. Theory , 58 (1989) , 68-77 .
5. Deeb , W. Multipliers And Isometries of Orlicz Spaces , in "Proceedings of the conference on Mathematical Analysis and its Applications , Kuwait Univ. Kuwait , Feb. 18-21" , 1985 .
6. Khalil , R. Best approximation in  $L^p(I,X)$  , Math. Proc. Cambridge Philos . soc. 94 (1983) , 277-279 .
7. Kreyszig , E. , Introductory Functional Analysis With Applications , John Wiley and sons , New York , 1978 .
8. Light , W. and Cheney , W. Approximation Theory In Tensor Product Spaces , Lecture Notes in Math. 1169 , springer-verlag , Berlin , 1985 .
9. Limaye , B.V. , Functional Analysis , Wiley Eastern limited , 1981 .
10. Rudin , W. , Real And Complex Analysis ; McGraw-Hill , New York , (1966) .

11. Siddiqi , A. H. , Functional Analysis With Applications , McGraw-Hill ,  
New York .
12. Singer , I. Best Approximation In Normed Linear Spaces By Elements  
of Linear Subspaces , springer-verlag , New York , Berlin , 1970 .